



Insecticidal Activity of Different Doses of *Acorus calamus* Essential Oil against *Sitophilus oryzae*

Mohammad Aasif Sheikh ^{a*}, Munazah Yaqoob ^b,
Fehim Jeelani Wani ^c, Tauseef A Bhat ^d,
Mudasir Gani ^b and Mohammad Anwar Bhat ^e

^a Division of Entomology, FoA, Wadura, SKUAST-K, Jammu and Kashmir, India.

^b Division of Entomology, FoH, Shalimar, SKUAST-K, Jammu and Kashmir, India.

^c Division of Agricultural Economics and Statistics, FoA, Wadura, SKUAST-K, Jammu and Kashmir, India.

^d Division of Agronomy, FoA, Wadura, SKUAST-K, Jammu and Kashmir, India.

^e Directorate of Extension, FoH, Shalimar, SKUAST-K, Jammu and Kashmir, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jsrr/2024/v30i62084>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113559>

Original Research Article

Received: 18/03/2024

Accepted: 20/05/2024

Published: 24/05/2024

ABSTRACT

The toxicity of different doses (3, 4, 5, 6 and 7 percent) of *Acorus calamus* essential oil, against insect pest *Sitophilus oryzae* were evaluated in the laboratory of Division of Entomology, FoA, Wadura, SKUAST-Kashmir during the year 2022. The various concentrations of *Acorus calamus* essential oil had significant effect on mortality of *Sitophilus Oryzae*. The highest mortality of 74.27

*Corresponding author: E-mail: aasifsheikh184@gmail.com;

Cite as: Sheikh, M. A., Yaqoob, M., Wani, F. J., Bhat, T. A., Gani, M., & Bhat, M. A. (2024). Insecticidal Activity of Different Doses of *Acorus calamus* Essential Oil against *Sitophilus oryzae*. *Journal of Scientific Research and Reports*, 30(6), 666–670. <https://doi.org/10.9734/jsrr/2024/v30i62084>

percent was recorded in 70 µl concentration of *Acorus calamus* essential oil on 12 Hours After Treatment (HAT) followed by 60 (71.01%), 50 (67.15%) and 40 (39.11%) µl concentration while untreated control recorded 5.13 percent mortality respectively. However, on 24 HAT the cumulative mortality increased to 76.35 in 70µl, 74.11 in 60 and 50 µl, 43.01 in 40 µl concentration. On 72 HAT the concentrations of 70, 60 and 50 µl recorded 94.11, 91.27 and 90.11 percent mortality respectively, while only 7.05 percent mortality was recorded in untreated control.

Keywords: *Acorus calamus*; essential oil; mortality; rice weevil; harmful species; seed viability.

1. INTRODUCTION

Insect pests that infest stored grain can lead to losses in weight, quality, commercial value, and seed viability. The majority of these pests, about seventy-five percent, belong to the Coleoptera order [1], with the most harmful species being found in the *Sitophilus* and *Tribolium* genera [2,3]. *Sitophilus oryzae* L. (Coleoptera: Curculionidae) is a widespread pest of significant economic impact that feeds internally by boring into stored grain. Adult weevils primarily consume the endosperm, thereby reducing the carbohydrate content, while the larvae prefer the germ, significantly depleting the grain's protein and vitamin levels. Insects that target the germ cause greater reductions in germination compared to others. Controlling arthropod pests in stored products has mainly involved the use of fumigants and residual chemical insecticides, in addition to maintaining proper hygiene practices [4,5]. The overuse of traditional chemical insecticides has led to several significant issues, such as insect resistance to these chemicals, the destruction of economically beneficial insects, environmental persistence, toxicity to humans and wildlife, and increased costs of crop production [6]. Numerous insects and mites have developed the ability to withstand nearly all pesticides used for their management due to cross-resistance and multiple resistance mechanisms [7]. Recognition of the deleterious effects of pesticides has spurred the exploration of alternative, less intrusive management approaches, such as employing the use of essential oils.

Numerous medicinal plants and spices have been utilized as pest control agents [8,9]. Farmers and researchers frequently report the effective use of plant materials for insect pest control, including ash [10,11], vegetable oils [12], plant extracts [13], and botanical powders [14]. It has been noted that certain plant-based preparations and traditional methods are much

safer than chemical insecticides [15], suggesting these materials should be explored for protecting stored products from pest infestations. *Acorus calamus* L. (Araceae), also known as "sweet flag," is a globally recognized ethnomedicinal and ethnobotanical plant. The rhizome of *Acorus calamus* exhibits a variety of pharmacological properties, including sedative, central nervous system depressant, behavior-modifying, anticonvulsant, memory-enhancing, anti-inflammatory, antioxidant, antispasmodic, cardiovascular, hypolipidemic, and immune-suppressive activities. *Acorus calamus* L. essential oil have also been used as fumigant against stored food products [16]. Keeping in view the insecticidal properties of essential oil of *Acorus calamus*, the present study was carried out to evaluate its various concentrations against rice weevil.

2. MATERIALS AND METHODS

2.1 Raising of Plant Material

The seedlings of *A. calamus* which were raised in polybags (Kharif, 2022) were procured from Regional cum Facilitation Centre North Zone II located at Faculty of Agriculture, Wadura Sopore and transplanted in the field with plant to plant and row to row distance of 15cm and 30cm, respectively. The rhizomes of mature plants were harvested, shade dried and ground to a fine powder and sieved through 30mm mesh. The powdered material was put to hydrodistillation for extraction of essential oil using Clevenger's apparatus. The essential oil of *A. calamus* thus obtained was collected in the vials and stored in refrigerator for further use.

2.2 Preparation of Different Concentrations of Essential Oil of *A. calamus*

Acetone was used as solvent for preparation of different concentrations. The doses 30, 40, 50,

60 and 70 µl corresponding to 3, 4, 5, 6 and 7 per cent were used for determining toxicity of *A. calamus* essential oil against *Sitophilus oryzae*.

2.3 Rearing of *Sitophilus oryzae*

The rearing of *S. oryzae* was carried in the laboratory of Division of Entomology, FoA, Wadura, SKUAST-Kashmir. The rice grains (K-39 Variety) infested with *S. oryzae*, was brought from different godowns/shops of Kashmir. Rice grains were taken in 5 litre capacity rearing jars and placed in hot air oven at 60°C for 24 hours for sterilization. The adult insects (both male and female) were collected randomly from the infested grains with the help of aspirator and transferred to the respective sterilized food placed in jars (5 litre capacity). The jars were covered with muslin cloth in order to supply adequate humidity to the grains and kept in B.O.D incubator at 28.5 ± 2°C and 65 ± 5% relative humidity to ensure proper egg laying and maintaining culture of rice weevil for further investigation. Mortality per cent was recorded after 12, 24, 36, 48, 60 and 72 hours after treatment using fumigation method. Test insects were considered dead if appendages do not move when prodded with a fine brush.

$$\text{Mortality (\%)} = \frac{\text{No. of insects dead}}{\text{Total No. of insects fumigated}} \times 100$$

2.4 Data Analysis

The data recorded in different treatments were subjected to analysis of variance (ANOVA) using R software.

3. RESULTS AND DISCUSSION

The insecticidal activity of Sweet flag against rice weevil is shown in Table 1. The highest mortality of 74.27 percent was recorded in 70 µl concentration of Sweet Flag on 12 Hours After Treatment (HAT) followed by 60 µl (71.01%), 50 µl (67.15%) and 40 µl (39.11%) concentration while untreated control recorded 5.13 percent mortality respectively (Table 1). However, on 24 HAT the cumulative adult mortality increased to 76.35 in 70 µl, 74.11 in 60 µl and 50 µl, 43.01 in 40 µl concentration (Table 1). On 72 HAT the concentrations of 70, 60 and 50 µl recorded 94.11, 91.27 and 90.11 percent mortality respectively, while only 7.05 percent mortality was recorded in untreated control.

It was observed that, the toxicity was dose dependent and enhanced with the increase in exposure time. Our results are in accordance with the findings of [17,18,19], who observed that concentration of *Acorus calamus* essential oil had significant effect on mortality of *Sitophilus Oryzae*.

Table 1. Effect of different concentrations of *Acorus calamus* essential oil against rice weevil

Concentration (µl/l of air)	Mortality (%) (HAT)						Mean
	12	24	36	48	60	72	
30	28.01 ^k (5.34)	33.03 ^{jk} (5.79)	39.11 ^{ijk} (6.29)	41.01 ^{ijk} (6.44)	42.13 ^{hij} (6.53)	43.05 ^{hij} (6.60)	37.72±2.4 2 (6.18)
40	39.11 ^{ijk} (6.29)	43.01 ^{ijk} (6.60)	45.03 ^{hij} (6.75)	46.13 ^{hi} (6.83)	47.11 ^{ghi} (6.90)	49.15 ^{fgh} (7.05)	44.92±1.4 3 (6.74)
50	67.15 ^{efg} (8.22)	74.11 ^{cdef} (8.64)	81.13 ^{abcde} (9.03)	88.03 ^{abc} (9.41)	89.01 ^{abc} (9.46)	90.11 ^{ab} (9.52)	81.59±3.8 0 (9.06)
60	71.01 ^{def} (8.46)	74.11 ^{cdef} (8.64)	80.03 ^{abcde} (8.97)	88.17 ^{abc} (9.42)	88.35 ^{abc} (9.43)	91.27 ^{ab} (9.58)	82.16±3.4 2 (9.09)
70	74.27 ^{cdef} (8.65)	76.35 ^{bcde} (8.77)	81.17 ^{abcde} (9.04)	86.03 ^{abcd} (9.30)	89.01 ^{abc} (9.46)	94.11 ^a (9.73)	83.49±3.1 1 (9.16)
Control	5.13 ⁱ (2.37)	5.05 ⁱ (2.36)	4.05 ⁱ (2.13)	6.01 ⁱ (2.55)	4.03 ⁱ (2.13)	7.05 ⁱ (2.75)	5.22±0.47 (2.39)
Mean	47.45±1.1 (6.92)	50.94±11.85 (7.17)	55.09±12.84 (7.46)	59.23±13.81 (7.73)	59.94±14.26 (7.77)	62.46±14.4 (7.93)	

HAT = Hours After treatment

* Values in parenthesis are square root transformed values

4. CONCLUSION

The study concludes that various concentrations of *Acorus calamus* essential oil had significant effect on mortality of *Sitophilus Oryzae* and the essential oil of *Acorus calamus* can be useful in controlling the stored product insect by surface treatment in godowns as an alternative to synthetic insecticides, further the efficacy of the botanical can be tested on other stored product insects also.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mohammad R, Roy S, Mishra BK, Khamari B. Biological repercussions of addition of botanicals to wheat grains on the progeny and damage potential of rice weevil (*Sitophilus oryzae* L.) at different temperature regimes. Cereal Research Communications. 2024, Mar; 52(1):203-13.
2. Khan AR, Selman BJ. On the mortality of *Tribolium castaneum* adults treated sublethally as larvae with pirimiphos methyl, Nosema whitei and pirimiphos methyl—*N. whitei* doses. Entomophaga. 1988, Sep;33:377-80.
3. Pinto Jr AR, Furiatti RS, Pereira PV, Lazzari FA. Avaliação de inseticidas no controle de *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae), e *Rhyzopertha dominica* (Fab.) (Coleoptera: Bostrichidae) em arroz armazenado. Anais da Sociedade Entomologica do Brasil. 1997;26:285-90.
4. Brooker DB, Bakker-Arkema FW, Hall CW. Drying and storage of grains and oil seeds. Springer Science & Business Media; 1992.
5. Adane K, Moore D, Archer SA. Preliminary studies on the use of *Beauveria bassiana* to control *Sitophilus zeamais* (Coleoptera: Curculionidae) in the laboratory. Journal of Stored Products Research. 1996, Apr 1;32(2):105-13.
6. Mario MB, Astuti LP, Hsu JL, Kafle L, Fernando I. Bioefficacy of eight different plant powders applied as fumigants against the adzuki bean weevil, *Callosobruchus chinensis*. Crop Protection. 2023, May 1;167:106200.
7. Metcalf RL. Changing role of insecticides in crop protection. Annual review of Entomology. 1980, Jan;25(1):219-56.
8. Lale NE. A laboratory study of the comparative toxicity of products from three spices to the maize weevil. Postharvest Biology and Technology. 1992;2(1):61-4.
9. Isman MB. Leads and prospects for the development of new botanical insecticides. Rev. Pestic. Toxicol. 1995;3:1-20.
10. Ofuya TI. Use of wood ash, dry chilli pepper fruits and onion scale leaves for reducing *Callosobruchus maculatus* (*Fabricius*) damage in cow-pea seeds during storage. The Journal of Agricultural Science. 1986, Oct;107(2):467-8.
11. Ajayi O, Arokoyo JT, Nezan JT, Olayinka OO, Ndirmbula BM, Kannike OA. Laboratory assessment of the efficacy of some local plant materials for the control of storage insect pests. Samaru Journal of Agricultural Research. 1987;5(1-2):81-6.
12. Aryal S, Poudel A, Bajracharya AS, Aryal LN, Kafle K. Toxicity evaluation of essential oil of clove (*Syzygium aromaticum*) bud against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae). Journal of Nepal Agricultural Research Council. 2023, Dec;31:22-35.
13. Chiasson H, Vincent C, Bostanian NJ. Insecticidal properties of a chenopodium-based botanical. Journal of Economic Entomology. 2004, Aug 1;97(4): 1378-83.
14. Gupta L, Srivastava M. Effect of *Withania somnifera* extracts on the mortality of *Callosobruchus chinensis* L. Journal of Biopesticides. 2008;1(2):190-2.
15. Yankanchi SR, Patil SR. Field efficacy of plant extracts on larval populations of *Plutella xylostella* L. and *Helicoverpa armigera* Hub. and their impact on cabbage infestation. Journal of Biopesticides. 2009;2(1):32-6.
16. Moshrefi Zenoozi Z, Soltaninezhad B, Hashemi M, Noori SM. A review of effective essential oils and their biologically active compounds to protect the safety of food stored against insect pests. Journal of Essential Oil Research. 2022;34(2):111-22.
17. Mansoor-ul-Hasan MS, Ullah E, Ahmad F, Wakil W. Insecticidal activity of different doses of *Acorus calamus* oil against

- Trogoderma granarium (Everts). Pak. J. Agri. sci. Val. 2006;43:1-2.
18. Abhijith N, Mohan S, Nelson SJ. Bioassay to detect the toxicity of sweet flag (*Acorus calamus*) 6 EC formulation against pests of stored spices viz., rusty grain beetle, *C. ferrugineus* (Stephens) cigarette beetle, *L. serricornis* (F.) and red flour beetle, *T. castaneum* (Herbst). Journal of Entomology and Zoology Studies. 2018;6(5):1921-1924.
19. Manjhu A, Lekha MK, Chhangani G, Kumar K. Host preference studies of rice weevil, *Sitophilus oryzae* L. on various cereals. 2022;11(1):1363-1367

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/113559>