



## **Antibacterial Activity of Bioflavonoid from Fruit Pulp of *Acacia nilotica* Willd**

**O. P. Oladosu<sup>1\*</sup>, N. R. Isu<sup>2</sup>, I. M. Aboh<sup>1</sup>, S. E. Okhale<sup>3</sup>, A. T. Orishadipe<sup>4</sup>  
and H. O. Egharevba<sup>3</sup>**

<sup>1</sup>Department of Microbiology and Biotechnology, National Institute for Pharmaceutical Research and Development, Abuja, Nigeria.

<sup>2</sup>Department of Microbiology, Faculty of Science, University of Abuja, Gwagwalada, Abuja, Nigeria.

<sup>3</sup>Department of Medicinal Plant Research and Traditional Medicine, National Institute for Pharmaceutical Research and Development, Abuja, Nigeria.

<sup>4</sup>Sheda Science Complex, Gwagwalada, Abuja, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. Author OPO responsible for drafting the proposal, plant collection antibacterial screening, analyzing the results and drafting of manuscript. Author NRI review the proposal and edited the reports. Author IMA antibacterial screening. Author SEO plant extraction and chromatography. Author ATO chemistry of the plant. Author HOE chemistry of the plant and drafting of manuscript. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/MRJI/2019/v28i430139

#### Editor(s):

(1) Dr. Marcin Lukaszewicz, Department of Biotransformation, Faculty of Biotechnology, University of Wroclaw, Wroclaw, Poland and Division of Chemistry and Technology Fuels, Wroclaw University of Technology, Wroclaw, Poland.

#### Reviewers:

(1) Debarshi Kar Mahapatra, Dadasaheb Balpande College of Pharmacy, India.

(2) Ronald Bartzatt, University of Nebraska, USA.

(3) Mustafa Sevindik, Akdeniz University, Turkey.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50244>

**Original Research Article**

**Received 04 May 2019  
Accepted 12 July 2019  
Published 01 August 2019**

### **ABSTRACT**

The emergence of multi-drug resistance in bacteria has led to call for research and development of new leads as antibiotics from medicinal plants. *Acacia nilotica* (Linn) is a plant of multipurpose medicinal uses, three bioactive flavonoids (methyl gallate, gallic acid and catechin) were isolated from its fruit pulps through a bioassay guided fractionation technique and characterized based on High Performance Liquid Chromatography, Liquid Chromatography-Mass Spectra and Nuclear Magnetic Resonance spectra. Antibacterial activity of these compounds was determined by

\*Corresponding author: E-mail: [petyem2001@yahoo.co.uk](mailto:petyem2001@yahoo.co.uk);

microplate tetrazolium dye assay of broth microdilution technique against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Escherichia coli*, and clinical isolates of *Salmonella typhi*, *Klebsiella pneumonia*, *Candida albicans* and *Bacillus subtilis*. Catechin, methyl gallate and gallic acid at 19.5, 39 and 39 µg/ml respectively caused a significant bio-reduction in cells of test organisms. Time kill kinetic study of the extract shows that there was percentage of growth reduction in test organisms at 2, 4, 6, 8 and 12 hrs of contact. The extent and rate of killing of the organism by the extract at 2 x MIC followed the same trend as rate of killing was time dependent. Antibacterial effects of these compounds are within the breakpoint of control drug chloramphenicol and could serve as leads in new drug development.

**Keywords:** *Acacia nilotica*; antibacterial; catechin; methyl gallate and gallic acid; bioactive compound; high through put techniques.

## 1. INTRODUCTION

*Acacia nilotica* Willd. Ex Delile is commonly known as Gum Arabic tree, Egyptian mimosa, Egyptian thorn, red thorn, Babool, babul in Indian [1]. It is recognized by the following names: Acacia, Acacia Arabica, Babhul – Hindi and Nepalese, Babool Baum – German etc. It has naturalized in several countries where it has been introduced as a medicinal, forage and fuel wood plant [2,1]. It is proverbial, medium sized tree and is broadly scattered in tropical and subtropical countries. It has an inspiring range of medicinal uses with potential antioxidant activity. It contains a profile of a variety of bioactive components such as gallic acid, (+)-catechin and methyl gallate. It also contains L-arabinose, catechol, galactan, galactoaraban, galactose, N-acetyl djenkolic acid, sulphoxides and pentosan. The seeds contain crude protein (18.6%), ether extract (4.4%), fiber (10.1%), nitrogen-free extract (61.2%), ash (5.7%) and silica (0.44%); phosphorus (0.29%) and calcium (0.90%) of dry matter (17%) [3]. Phytochemical analysis of stem bark of *A. nilotica* showed the presence of carbohydrates, saponins, tannins, and cardiac glycosides [4].

As a multipurpose medicinal plants, leaves, flowers, seeds, roots, fruits and gum act as anti-tuberculosis [5], anti-hypertensive and anti-spasmodic, antibacterial and antifungal [6], antiplasmodial [7], and antioxidant [8,9,10]. Toxicologically, *A. nilotica*, at 2 and 8% levels, has a low toxicity potential [11]. In a survey of potentially allergenic plants in Pondicherry, it was reported likely to cause pollen allergy [12]. *A. nilotica* has a wealth of medicinal uses for stomach upset and pain, the bark is chewed to protect against scurvy, an infusion is taken as therapy for dysentery and diarrhea [13]. In Nigeria, it is one of the customary drugs for treating diarrhea and it has been authenticated to

have an antidiarrheal property [3] and antimicrobial activity against neuro-pathogenic *E. coli* K1, MRSA and *Klebsiella pneumonia* [14].

This study describes the inhibitory effect of bioactive molecules of *A. nilotica* as leads for drug development against pathogenic organisms using tetrazolium salts reduction as indicators of cell inhibition.

## 2. METHODOLOGY

### 2.1 Chemicals/Reagents/Equipment

Tetrazolium dye, Nutrient broth No 1, Muller Hinton Agar from Fluka Analytical, Sigma-Aldrich Co., USA; Nutrient Agar from HIMedia Laboratories Pvt Ltd., India; chloramphenicol discs (REF OD 278R, HIMedia Laboratories Pvt. Ltd, India), Uv/vis spectrophotometer (Jenway 6405, Britain), nuclear magnetic resonance (300 MHz, Varian machine), HPLC and LCMS.

### 2.2 Extract Preparation and Isolation

The pulp of *A. nilotica* fruit was collected from Suleja, Niger state, Nigeria, identified at Herbarium Unit of National Institute for Pharmaceutical Research and Development, Abuja, Nigeria and herbarium specimen was deposited. This part was air dried to constant weight at room temperature and pulverized. Extraction of both polar and non-polar components of the plants using 70% aqueous methanol was carried out. One hundred grams of dried plant part was soaked in 70% aqueous methanol (1000 ml) for 24 hrs. The extract was filtered, concentrated on rotary evaporator at 40°C and freeze dried. A portion of the crude extract was used to determine the preliminary antibacterial effect of the plant, while the other portion was subjected to bioassay guided isolation of the bioactive molecules (Fig. 1).

Fractionation of the crude extract of *A. nilotica* (seeds) was carried out according to Orishadipe et al. [15]. Activated silica gel (50 g) was packed by a wet method into a column. The extract (2.6g) which was absorbed on silica gel (60 – 120 mesh) and dried was loaded on the column. Gradient elution was performed with 100 ml of each mobile phase mixture in a series. The elution was performed in a flash chromatography. The mobile phase consisted of hexane, ethyl acetate, methanol and water, starting from 100% hexane and 10% increment in the next polar component. The final elution was performed with 70% methanol in water until the column appeared exhausted with a sign of colourless silica gel. The eluates were monitored by thin layer chromatography (TLC) using normal phase pre-coated silica gel K5 TLC plates. The TLC mobile phase consists of a mixture of ethyl acetate: hexane (7:3). The eluates were combined based on the similarity of TLC fingerprint to give six fractions labelled as FR1-FR6. The pooled fractions (0.8 g) were absorbed on Merck – Kiesegel and introduced into column packed with the same absorbent. It was eluted with petroleum ether and an increasing gradient of ethyl acetate. A total of 6 fractions of 100 ml each were collected. Fractions with same similarity (2-4) were pooled together into 3 portions viz; P1, P2 and P3 based on their TLC

profile (Fig. 2). On staining with ferric chloride solution showed blue black colouration which is an indication of phenolic compounds. These fractions were purified further by HPLC technique by separating on a reverse phase column, eluting with a gradient of acetonitrile 5% - 98% in water for a period of 40 minutes. Some 100 mg of extract was dissolved in 1 ml of methanol, and water was added gradually to the point of precipitation before injecting on the HPLC reversed phase (C-18) column. Fractions were collected at intervals of one minute. The structural elucidation of the compounds was done using Nuclear Magnetic Resonance (300 MHz, Varian machine) and LC-MS techniques. [2008]. The pure fractions from TLC analysis were dissolved in deuterated methanol (CD<sub>3</sub>OD) and transferred to NMR tube for measurement of proton (<sup>1</sup>H) and <sup>13</sup>C (carbon 13). Masses were acquired using liquid chromatography- Mass Spec (LC-MS).

### 2.3 Phytochemical Analysis

Phytochemical analysis of fruit pulp of *A. nilotica* was carried out according to Trease and Evans [16]. Metabolites tested for include alkaloid, saponins, tannins, anthraquinone, cardiac glycosides and flavonoids.




Fig. 1. Extraction procedure of fruit pulps of *A. nilotica*

## 2.4 Antibacterial Activity of the Crude Extract

### 2.4.1 Extract solution preparation

The extract was dissolved in dimethylsulphoxide (DMSO) and further diluted to 40 mg/ml concentration. The extract was sterile-filtered with 0.22 µm syringe filter (Fisher brand). From this, a 2 mg/ml concentration of extract in agar was prepared by dispensing 1ml of the extract solution (40 mg/ml) into 19 ml of molten Mueller Hinton Agar, maintained in water bath at 45°C. The molten agar/extract was poured into sterile Petri dish, allowed to gel and dried to remove moisture. Overnight broth cultures of the test organisms viz; *S. aureus* (ATCC 28923), *P. aeruginosa* (ATCC 27853), *E. coli* (ATCC 25922) and clinical isolates of *S. typhi*, *K. pneumonia*, *C. albicans* and *B. subtilis* were standardized by diluting to 10<sup>7</sup> cfu/ml using a uv/vis-spectrophotometer as described by Dominguez et al. [17]. Two to three colonies of 20 hr growth on Mueller Hinton agar of the organisms to be studied were suspended on 50 ml pre-warmed (37°C) Mueller Hinton broth. The suspension was incubated overnight at 37° C, diluted 1/2500 in the same pre-warmed medium and incubated in water bath with agitation (50 rpm). The optical density of the culture was monitored at 450 nm until absorbance of 0.1 was reached (equivalent 2.5-3.0 x 10<sup>7</sup> cfu/ml for *E. coli* and *P. aeruginosa*, 1.8-2.0 x 10<sup>7</sup> for *S. aureus* and *B. subtilis*). The agar containing extract was inoculated by dropping 50 µl of standardized organism on the surface of the agar, allowed to stand for 2 hrs. The plate was incubated for 24 hrs at 37°C. The experiment was done in duplicate. Control plates containing 30 µg/ml concentration of chloramphenicol (prepared by dissolving 30 mg of chloramphenicol powder in 10 ml 10% dimethyl sulfoxide (DMSO) and further diluted by taking 10 µl solution in 10 mL nutrient broth) was used as drug control to confirm the susceptibility of the test organisms. Agar containing solution of the extract served as extract sterility control. Post incubation, the presence or absence of growth of colonies of test organisms on agar plates indicated positive or negative results of the crude extract.

### 2.4.2 Minimum inhibitory concentration of molecules from *Acacia nilotica* (P1, P2 and P3)

The minimum inhibitory concentration of molecules of *A. nilotica* was determined by broth microdilution method according to British Society

of Antimicrobial Susceptibility [18] recommendation with a slight modification using tetrazolium colorimetric assay as growth/inhibition indicator [19,20]. Ten milligrams of each molecule was dissolved in Muller Hinton broth. 100 µl of each of the solutions was transferred to the first column of micro well plate, from where 50 µl was transferred to 2<sup>nd</sup> well already containing 50 µl Muller Hinton broth, mixed thoroughly and the procedure was repeated through to well 11 where 50 µl was discarded. All the wells were inoculated with 50 µl of overnight diluted cultures of each of *S. aureus*, *Pseudomonas aeruginosa*, *E. coli*, *S. typhi*, *K. pneumonia*, *C. albicans* and *B. subtilis* prepared as described above. The plates were incubated for 24 hrs at 37°C. Post incubation the plates were stained with tetrazolium dye; 3-(4, 5-dimethylthiazol-2-yl)-2, 5 diphenyltetrazolium (MTT). MICs were defined as lowest concentration of antimicrobial agents that red formazan of MTT was not observed.

### 2.4.3 Time kill Kinetic antibacterial study of fruit extract of *Acacia nilotica*

One hundred milligram per ml concentration of the crude extract was prepared in sterile water and diluted to 2 mg/ml concentration (being 2 X mic of the crude extract) filtered by centrifuging for 10 mins at 4500 rpm to remove woody sediments. The filtrate was inoculated with 10<sup>5</sup> cfu/ml of *S. aureus* (ATCC 28923) and *E. coli* (ATCC 25922) and incubated at 37°C, percentage growth decrease was measured at 2 hrs interval. Exactly 0.5 ml of each suspension was withdrawn at the appropriate time intervals and transferred to 4.5 mL nutrient broth recovery medium containing 3% Tween 80 to neutralize the carry-over effects of the antimicrobial compounds from the test suspensions. The suspension was shaken properly then serially diluted in sterile physiological saline. Exactly 0.5 mL of the final dilution of the test organism was transferred into pre-sterile Nutrient agar at 45°C and plated out. The plates were allowed to set and incubated upside down at 37°C for 72 h. Optical density of growing culture in extract solution was measured using uv-spectrophotometer (Jenway 6405 uv/vis, UK) at 520 nm to determine decrease in cells growth. Control experiment which was set up without the inclusion of antimicrobial agent (i.e. excluding the crude extract). Viable counts were made in triplicates for each sample. Depression in the viable counts indicated killing by the antimicrobial agent. Statistical analyses were performed with SPSS version 16.0 (SPSS Inc., Chicago, IL,

USA) software. The level of significance for all statistical tests was set at  $p < 0.05$ .

### 3. RESULTS AND DISCUSSION

Phytochemical analysis of fruit pulp of *A. nilotica* showed that the plant contains alkaloid, saponins, tannins (phlobatannins), anthraquinone and flavonoids. The presence of these phytochemicals could be responsible for the observed activity. For instance, tannins therapeutically have antiseptic properties and their precipitating quality is used in detecting gelatin, proteins and alkaloids [21]. Flavonoids and phenolic compounds are plants metabolites with at least one hydroxyl group [22] and are often found effective *in vitro* as antimicrobial substance against a wide array of microorganisms [23]. The presence of these phytochemicals could be responsible for the antibacterial activity recorded in this study.

Bioassay guided fractionation of the crude extract of fruit pulp of *A. nilotica* yielded 3 distinct spots on TLC (Fig. 2). The compounds were analyzed on Liquid Chromatography- Mass Spectrometer (LCMS) for purity and mass. Three compounds were isolated from the fractions by the preparative. The pure compounds were fractions 5, 9 and 12. Fractions in between were mixtures of these compounds as indicated by NMR spectra.

The <sup>1</sup>H NMR of fractions 5 revealed it to be methyl gallate, 9 to be gallic acid and 12 to be catechin through a library search [24,15].

#### 3.1 Characterization of Isolated Compounds

The first compound P1 (fraction 5) was a white solid with a melting point of 202-204 °C (Lit. 201-204 °C). The <sup>1</sup>H NMR revealed the presence of aromatic protons at  $\delta$ 7.2, which integrated for two protons at positions 2 and 6 and that of methoxyl at  $\delta$ 3.3 which integrated for 3 protons at position 8. This indicated a simple aromatic system. The broad band was decoupled as; <sup>1</sup>H NMR: 7.20 (2H, H-2 and H-6), 5.0 (OH), 3.3 (3H, CH<sub>3</sub>), <sup>13</sup>C NMR: 167.9 (C-7), 146.6 (C-3), 146.6 (C-5), 139.4 (C-4), 122.2 (C-1), 110.3 (C-2), 110.3 (C-6), 52.5 (OMe). <sup>13</sup>C NMR revealed the presence of 8 carbons at  $\delta$  (167.9 for one C=O, 110.3 for two CH, 52.5 for OCH and four quaternary aromatic carbons at 122.2, 138.4, 146.6, 146.6) (Table 1). HRMS of this fraction gave 184.03717 amu, which was consistent with molecular formula C<sub>8</sub>H<sub>8</sub>O<sub>5</sub>. A library search revealed it to be methyl gallate (methyl 3, 4, 5-trihydroxybenzoate) [15,24]. The sample was also compared with an authentic sample by TLC and was confirmed to be methyl gallate: Methyl 3, 4, 5-trihydroxybenzoate (Fig. 3). Molecular formula; C<sub>8</sub>H<sub>8</sub>O<sub>5</sub>.





Fig. 2. Thin layer chromatography plate of active fraction of *A. nilotica* (showing 3 distinct spots P1, P2 and P3)



**Fig. 3. Chemical structure of methyl gallate with molecular formula C<sub>8</sub>H<sub>8</sub>O<sub>5</sub>**


**Table 1. Carbon 13 and proton NMR spectra of methyl gallate (P1) and gallic acid (P2)**

| Position | Group            | Compound P1                                      |   | Compound P2                                      |   |
|----------|------------------|--|---|--|---|
|          |                  | <sup>13</sup> C Chemical Shift (δ <sub>C</sub> ) | Proton chemical shift (δ <sub>H</sub> ) | <sup>13</sup> C chemical shift (δ <sub>C</sub> ) | Proton chemical shift (δ <sub>H</sub> ) |
| 1        | C                | 122.2  | -                                       | 122.2  | -                                       |
| 2        | CH               | 110.3  | 7.2                                     | 110.3  | 7.0                                     |
| 3        | C-OH             | 146.6  | 6.8                                     | 146.6  | 4.789                                   |
| 4        | C-OH             | 139.4  | 5.0                                     | 140.0  | 4.789                                   |
| 5        | C-OH             | 146.6  | 6.8                                     | 146.6  | 4.789                                   |
| 6        | CH               | 110.3  | 7.2                                     | 110.3  | 7.0                                     |
| 7        | C=O              | 167.9  | -                                       | -  | -                                       |
| 8        | OCH <sub>3</sub> | 52.5   | 3.30                                    | -  | -                                       |

The second compound (P2) was a solid, light brownish in colour, m. p. = 237-239°C. <sup>1</sup>H NMR: 7.0 (2H, H-2 and H-6), 4.789 (OH), <sup>13</sup>C NMR: 170 (C-7), 146.6 (C-3), 146.6 (C-5), 139.4 (C-4), 122.2 (C-1), 110.3 (C-2), 110.3 (C-6). The <sup>1</sup>H NMR of this compound revealed the presence of aromatic protons at δ 7.0, which integrated for two protons at positions 2 and 6 and the OH proton at position 7 at δ 3.212. The broad band decoupled (Fig. 4.) <sup>13</sup>C NMR revealed a total of 7 carbons at δ (170 for one C=O, 110.3 for two CH, and 4 quaternary aromatic carbons at 122.2, 138.4, 146.6, 146.6). These were almost identical with that of methyl gallate except for the absence of the OCH<sub>3</sub> <sup>13</sup>C NMR peak. As a result

this compound was subsequently identified as gallic acid [15]. Fig. 4.


<sup>1</sup>H-NMR spectra of the third compound (P3) showed peak at δ 4.56 (H-2, d), 4.01 (H-3, ddd), 2.54 (H-4, dd), 2.90 (H-4, dd), 5.87 (H-6, d), 6.01 (H-8, d) 6.89 (H-2', d), 6.79 (H-5', d), 6.73 (H-6', dd) and 8.00 (phenolic protons, m). <sup>13</sup>C-NMR, Carbon atoms showed peaks at δ TMS 80.9 (C-2), 66.3 (C-3), 27.7 (C-4), 93.9 (C-6), 95.1 (C-8), 114.5 (C-2'), 115.1 (C-5'), 18.4 (C-6') (Table 2). The NMR chemical shifts correlate well with those available in literature for catechin [25; 26]. The compound was identified as catechin (Fig. 5).



**Fig. 4. Chemical structure of gallic acid with molecular formula C<sub>7</sub>H<sub>6</sub>O<sub>5</sub>**

Table 2. Carbon 13 and proton NMR spectra of catechin (P3)

| Position | Carbon 13 chemical shift ( $\delta_C$ ) | Proton chemical shift ( $\delta_H$ ) |
|----------|---|--------------------------------------|
| 1        | -                                       |                                      |
| 2        | 80.9                                    | 4.56                                 |
| 3        | 66.3                                    | 4.00                                 |
| 4        | 27.7                                    | 5.58                                 |
| 5        | 156.4                                   |                                      |
| 6        | 93.9                                    |                                      |
| 7        |   |                                      |
| 8        | 95.1                                    |                                      |
| 9        |   |                                      |
| 1'       |   |                                      |
| 2'       | 114                                     | 6.89                                 |
| 3'       |   |                                      |
| 4'       |   |                                      |
| 5'       | 115.1                                   | 6.79                                 |
| 6'       | 18.4                                    | 18.4                                 |



Catechin

Fig. 5. Chemical structure of catechin with molecular formula  $C_{15}H_{14}O_6$ 

### 3.2 Antibacterial Activity

Antibacterial activity of the crude fractions and compounds from fruit pulps of *A. nilotica* are as shown in Table 3. The result shows that the crude extract was significantly active against all the test organisms but at varying degree of concentration. Minimum inhibitory concentration against *B. subtilis* and *S. aureus* were 500 and  $750 \pm 0.0$   $\mu\text{g/ml}$  respectively while the susceptibility of Gram negative organisms (*E. coli*, *P. aeruginosa*, *S. typhi*, *K. pneumonia*) and *C. albicans* were slightly higher at 1000 – 2000  $\mu\text{g/ml}$  respectively). These concentrations are higher than the mic breakpoint for chloramphenicol ( $30 \pm 0.1$   $\mu\text{g/ml}$ ) the control drug. Hence the need for further purification of the extract to compound level for better activity. The fractions (F1-F6) produced by column chromatography with TLC finger printing yielded better minimum inhibitory concentrations (MICs) at 78 –  $156 \pm 0.0$   $\mu\text{g/ml}$  respectively. Fraction F1 had no antibacterial effect, fraction F5 had mic of 156  $\mu\text{g/ml}$  against all the test organisms while F6 was only active against *S. aureus* and *B. subtilis*

at 156  $\mu\text{g/ml}$ . Fractions F2-F4 produced the best antibacterial effects at 78  $\mu\text{g/ml}$  against Gram positive organisms and 156  $\mu\text{g/ml}$  against Gram negative organisms. These fractions have a lot in common; TLC profile, positive test for phenolic and antibacterial effects. Microplate tetrazolium assay of isolated molecules against the pathogenic organisms as shown in Table 1, showed that methyl gallate, gallic acid at 39  $\mu\text{g/ml}$  and catechin at 19.5  $\mu\text{g/ml}$  concentration are significantly ( $p < 0.05$ ) more effective against *E. coli*, *S. aureus* and *B. subtilis* than against *P. aeruginosa*, *S. typhi* and *K. pneumonia* (78  $\mu\text{g/ml}$ ). Catechin has shown to be the most effective of the molecules with MIC of  $19 \pm 0.5$   $\mu\text{g/ml}$  against *S. aureus* and *B. subtilis* and 39  $\mu\text{g/ml}$  against other Gram negative organisms.

This result shows that the purer the fractions the better the antibacterial activity. The use of microplate tetrazolium assays to measure cell proliferation has increased exponentially since their introduction [19]. Nevertheless, these assays do not actually measure the number of viable cells in a culture or their growth but rather,

**Table 3. Antibacterial activity of crude, fractions and compounds of fruit pulp of *A. nilotica***

| Molecules µg/ml | Microorganisms   |                      |                |                    |                 |                     |                    |
|-----------------|------------------|----------------------|----------------|--------------------|-----------------|---------------------|--------------------|
|                 | <i>S. aureus</i> | <i>P. aeruginosa</i> | <i>E. coli</i> | <i>B. subtilis</i> | <i>S. typhi</i> | <i>K. pneumonia</i> | <i>C. albicans</i> |
| Crude extract   | 750 ± 0.0        | 1000 ± 0.0           | 1000 ± 0.0     | 500 ± 0.0          | 2000±0.0        | 2000±0.0            | 1000±0.0           |
| F1              | NA               | NA                   | NA             | NA                 | NA              | NA                  | NA                 |
| F2              | 78               | 156                  | 156            | 78                 | 156             | 156                 | 156                |
| F3              | 78               | 156                  | 156            | 78                 | 156             | 156                 | 156                |
| F4              | 78               | 156                  | 156            | 78                 | 156             | 156                 | 156                |
| F5              | 156              | 156                  | 156            | 156                | 156             | 156                 | 156                |
| F6              | 156              | NA                   | NA             | 156                | NA              | NA                  | 156                |
| Methyl gallate  | 39 ± 0.0         | *78 ± 0.0            | 78 ± 0.0       | 39 ± 0.0           | 78              | 78                  | 39                 |
| Gallic acid     | 39 ± 1.0         | 78 ± 0.0             | 78 ± 0.0       | 39 ± 0.0           | 78±0.0          | 78±0.0              | 39±0.0             |
| Catechin        | 19.5 ± 0.0       | 39 ± 0.0             | 39 ± 0.0       | 19.5 ± 0.0         | 39±0            | 39±0.0              | 39±0.0             |
| Chloramphenicol | 30± 0.0          | 30 ± 0.00            | 30 ± 0.0       | 30±0.0             | 30±0.0          | 30±0.0              | 30±0.0             |

Key: F= Fractions, NA = No activity, no growth inhibition



an integrated set of enzyme activities that are related in various ways to cell metabolism. They utilize the cofactor nicotinamide adenine dinucleotide (NADH) and the dye 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium (MTT), and other substrates like succinate and pyruvate which may also contribute to their reduction. Depending on the particular dye chosen, reduction will be linked in various ways to cofactor/substrate production, utilization and compartmentalization, and can be associated with the plasma membrane, intracellular membranes, organelles and cytosol. Reduction can vary widely within and between cell populations depending on the cell growth conditions, phase of cell growth and stage of the cell cycle [20].

In the case of antibacterial activity of methyl gallate, the result is in agreement with a similar study [27] that methyl gallate from the leaves of *Toona sureni* had antibacterial activity against *E. coli*, *S. aureus* and *B. subtilis*. Other similar studies have reported that gallic acid by mechanism of action in *P. aeruginosa*, *E. coli*, *S. aureus* and *Listeria monocytogens* led to irreversible changes in membrane properties (charge, intra and extracellular permeability, and physicochemical properties) through hydrophobicity changes, decrease of negative surface charge, and occurrence of local rupture or pore formation in the cell membranes with consequent leakage of essential intracellular constituents [28]. Antibacterial activity of gallic

acid of *A. nilotica* in this study corroborate the report that gallic acid inhibited the growth of *Campylobacter jejuni* and *E. coli* strains at 15.63-250 µg/ml. [12].

Prolonged incubation of the microplates was used to determine the end point assay of the molecules as bacteriocidal. Wells with bacteriostatic effect at concentrations lower than 78 µg /ml showed tetrazolium reduction after 48 hours of incubation while bacteriocidal concentrations of 78 µg /ml upward remained colourless. This assay has been used to distinguish between dormant and metabolically active microbial cells (Tab. 3). The test organisms were able to reduce tetrazolium dyes in their electron transport chain, generating results within hours.

### 3.3 Time Kill Kinetic Antibacterial of Crude Extract

The time kill kinetic antibacterial of crude extract against *S. aureus* and *E. coli* are as shown in Figs. 6 and 7. The percentage of growth reduction in *S. aureus* at 2, 4, 6, 8 and 12 hrs of contact were 52.2, 60.8, 70, 80 and 99.2% respectively. Percentages of *E.coli* growth reduction for each corresponding time were; 45.7, 51.7, 64.16 and 99.9% respectively. The extent and rate of killing of the organism by the extract at 2 x MIC followed the same trend as the concentrations of the extract increased with increase in contact time, the percentage of the organisms killed also increased.




Fig. 6. Effect of extract of *A. nilotica* on the growth of *S. aureus* at different time




Fig. 7. Effect of extract of *A. nilotica* on the growth of *E. coli* at different time

#### 4. CONCLUSION

In conclusion, this study has demonstrated the antibacterial properties of bioflavonoids from fruit pulps of *A. nilotica*. Step by step bioassay guided isolation shows an improved antibacterial effect from crude extract to pure compounds. The results of the minimum inhibitory concentrations of the compounds showed that the compounds are active against all test organisms, as seen the isolated compounds have antibacterial effect close to the MIC breakpoint of the control drug, although methyl gallate and gallic acid have better activity than the crude extract, catechin has shown to be the most active compound against all the test organisms. The time kill antibacterial study was able to demonstrate time dependent effect of the crude extract against the test organisms. Generally, the compounds isolated could serve as leads in the search for new and potent antibacterial compounds from indigenous medicinal plants.

#### ACKNOWLEDGEMENT

The authors are grateful to Nigerian Conservative Foundation Lekki, Lagos State, Nigeria for Chief SL Edu memorial research grant.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Bennison JJ, Paterson RT. The use of Trees by Livestock. Acacia Production Programme. 1994;1:160-164.
- Atif Ali, Akhtar Naveed, Barkat Ali Khan, Muhammad Shoaib Khan, Akhtar Rasul Shahiq-UZ-Zaman, Nayab Khalid, Khalid Waseem, Tariq Mahmmod and Liaqat Ali. Acacia nilotica: A plant of multipurpose medicinal uses. Journal of Medicinal plants Research. 2012;6(9):1492-1496.
- Ogbadoyi EO, Garba MH, Kabiru AY, Mann A, Okogun J. Therapeutic evaluation of Acacia nilotica (Linn) stem bark extract in experimental African trypanosomiasis. International Journal of Applied Research of Natural Products. 2011;4(2):11-18.
- Sawant RS, Godghate AG, Sankpal SA, Walaki SA, Kankanwadi SS. Phytochemical analysis of bark of Acacia nilotica. Asian Journal of Plant Science and Research (Pelagia Research Library). 2014;4(2):22-24.B.
- Oladosu P, Isu NR, Ibrahim K, Okolo P and Oladepo DK. Time kill-kinetics antibacterial study of *Acacia nilotica*. African Journal of Microbiology Research. 2013b;7(46):5248-5252.
- Banso A. Phytochemical and antibacterial investigation of bark extracts of Acacia nilotica. Journal of Medicinal Plants Research. 2009;3:082-085.

7. Jigam AA, Akanya HO, Dauda BEN, Okogun JO. Polygalloytannin is effective against *Plasmodium berghei* in mice. *J. Med. Plants Res.* 2010;4(12):1169-1175
8. Del WE. *In vitro* evaluation of peroxyl radical scavenging capacity of water extract/fractions of *A. nilotica* (L). *African Journal of Biotechnology.* 2009;8(7):1270-1272.
9. Amos S, Akah PA, Odukwe CJ, Gamaniel KS, Wambebe C. The pharmacological effect of an aqueous extract from *Acacia nilotica* seeds. *Phytotherapy Research.* 1999;13:683-685
10. Kalavaini T, Mathew L. Free radical scavenging activity from leaves of *Acacia nilotica* (L) Will. ex. Delile, an Indian medicinal tree. *Food Chem. Toxicol.* 2010;48:298-305.
11. Al-Mustafa ZH, Dafallah AA. A study on the toxicology of *Acacia nilotica*. *Am. J. China Med.* 2000;28(1):123-129.
12. Amrreta Sarjit, Yi Wang, Gary A. Dykes. Antimicrobial activity of gallic acid against thermophilic *Campylobacter* is strain specific and associated with a loss of calcium ions. *Food Microbiology.* 2014;46, 227-233.
13. Misar A, Bhagat R, Mujumdar AM. Antidiarrhoeal activity of *Acacia nilotica* Willd. bark methanol extract. *Hindustan Antibiot Bull.* 2008;49-50(1-4):14-20.
14. Saba R, Muhammad F, Shahida H and Naveed Ak. Antibacterial and Cytotoxicity of *Acacia nilotica* Lam (Mimosaceae) Methanol extracts against extended spectrum Beta- lactamase producing *Escherichia coli* and *Klebsiella* species. *Trop. J. Pharm. Res.* 2011;10(6):785-791
15. Orishadipe AT, Oladosu P, D'Ambroso M, Adesomoju AA, Ibrahim K, Barry CE 111 and Okogun JI. Isolation and characterization of 1, 2, 3, 4, 6-pentagalloyl glucose an antimycobacterial agent from the leaves of *Entandrophragma angolense*. *Journal of Phytomedicine and Therapeutics.* 2008;13:16–22.
16. Trease GE, Evans WC. *Pharmacognosy* 15th Edn., WB Sanders, London. 2002;406. [ISBN: 8131200876]
17. Dominguez MC, Manuel de la Rosa, Victoria MB. Application of spectrophotometric method for the determination of post- antibiotic effect and comparism with viable counts in agar. *Journal of Antimicrobial Chemotherapy.* 2001;47:391–398.
18. British Society of Antimicrobial Susceptibility. Guidelines for the laboratory diagnosis and susceptibility testing of methicillin-resistant *Staphylococcus aureus* (MRSA). *Journal of antimicrobial chemotherapy.* 2005;56:1000-1018.
19. Mozmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxic assays. *Journal immunology Methods.* 1983;65:55-63.
20. Marshall NJ, Goodwin CT and Holt SJ. A critical assessment of micro culture tetrazolium assays to measure cell growth and function. *Growth Regulation.* 1995;5; 69-84
21. Adamczyk J, Simon V, Kitunen S, Adamczyk A, Smolander. Tannins and their complex interaction with different organic nitrogen compounds and enzymes: Old paradigms versus recent advances. *Chemistry Open (Reviews)* 2017;6:610– 614. DOI: 10.1002/open.201700113
22. Duangjai Tungmunnithum, Areeya Thongboonyou, Apina Pholboon and Aujana Yangsabai. Flavonoids and other phenolic compounds from medicinal plants for pharmaceutical and medical aspects: An overview. *Medicines (Basel).* 2018;5(3): 93. Doi: 10.3390/medicines 5030093
23. Bennett RN, Wallsgrove RM. Secondary metabolites in plant defense mechanisms. *Phytotherapy Research.* 1994;127:617-633.
24. Pouchet J, Charles Jacquelyn Benlinke. *The Aldrich library of 1H and 13CNMR (FT- Spectra).* Aldrich Chemical Co. Inc. 1993;111.
25. Hye MA, Taher MA, Ali MY, Ali MU, Shahed Zaman. Isolation of (+) catechin from *Acacia catechu* (Cutch tree) by Convenient Method. *J. Sci. Res.* 2009;1(2):300-305.
26. Kiehlmann E, Tracey AS. Proton magnetic resonance spectra of catechin and bromocatechin derivatives: C6' vs. C8-substitution. *Can. J. Chem.* 1986;64: 1998-2005.

27. Taufik M. Ekaprasada, Hazli Nurdin, Sanusi Ibrahim, Dachriyanus. Antibacterial Activity of Methyl Gallate Isolated from the Leaves of *Toona sureni*. International Journal on advanced Science Engineering Information Technology; 2005.
28. Borges A, Ferreira C, Saavedra MJ, Simões M. Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. Microbial Drug Resistant. 2013;(4): 256-65.

© 2019 Oladosu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/50244>*