



A Note on Sum Formulae of Generalized Pentanacci Sequence

Yüksel Soykan ^a and Erkan Taşdemir ^{b*}

^aDepartment of Mathematics, Art and Science Faculty, Zonguldak Bülent Ecevit University, 67100, Zonguldak, Turkey.

^bPınarhisar Vocational School, Kırklareli University, 39300, Kırklareli, Turkey.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/arjom/2024/v20i10854>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124484>

Received: 03/08/2024

Accepted: 07/10/2024

Published: 22/10/2024

Original Research Article

Abstract

During this paper, we first demonstrate the closed forms of sum formulae both $\sum_{\lambda=0}^n \lambda z^\lambda A_\lambda$ and $\sum_{\lambda=1}^n \lambda z^\lambda A_{-\lambda}$ for the generalized Pentanacci numbers. Then, we grant summation formulae for the sequences such as Pentanacci, Pentanacci-Lucas and other fifth-order iteration sequences.

Keywords: Pentanacci numbers; pentanacci-lucas numbers; sum formulae; summing formulae.

2010 Mathematics Subject Classification: 11B37, 11B39, 11B83.

*Corresponding author: E-mail:erkantasdemiir@hotmail.com;

Cite as: Soykan, Yüksel, and Erkan Taşdemir. 2024. "A Note on Sum Formulae of Generalized Pentanacci Sequence". Asian Research Journal of Mathematics 20 (10):184-204. <https://doi.org/10.9734/arjom/2024/v20i10854>.

1 Introduction

First of all, we revisit the Pentanacci sequences. The Pentanacci sequence is a higher-order generalization of the well-known Fibonacci sequence. Defined by a linear recurrence relation, each term in the Pentanacci sequence is the sum of the five preceding terms:

$$A_\eta = A_{\eta-1} + A_{\eta-2} + A_{\eta-3} + A_{\eta-4} + A_{\eta-5}, \eta \geq 5;$$

with initial conditions A_0, A_1, A_2, A_3, A_4 set to predefined values. The Pentanacci sequence has applications in theoretical mathematics, particularly in combinatorics and the analysis of recursive algorithms. Its study reveals deeper insights into the properties of linear recursions and their behavior in higher dimensions, offering attractive connections to both discrete mathematics and dynamical systems.

Now, we recall the generalized Pentanacci numbers. The generalized Pentanacci sequence $\{A_\eta(A_0, A_1, A_2, A_3, A_4; r, s, t, u, v)\}_{\eta \geq 0}$ (or $\{A_\eta\}_{\eta \geq 0}$) is defined in the following way:

$$\begin{aligned} A_\eta &= rA_{\eta-1} + sA_{\eta-2} + tA_{\eta-3} + uA_{\eta-4} + vA_{\eta-5}, \\ A_0 &= c_0, A_1 = c_1, A_2 = c_2, A_3 = c_3, A_4 = c_4, \eta \geq 5, \end{aligned} \tag{1.1}$$

where A_0, A_1, A_2, A_3, A_4 are arbitrary numbers (real or complex) and r, s, t, u, v are real numbers. The sequence $\{A_\eta\}_{\eta \geq 0}$ have negative subscripts by definition as

$$A_{-\eta} = -\frac{u}{v}A_{-\eta+1} - \frac{t}{v}A_{-\eta+2} - \frac{s}{v}A_{-\eta+3} - \frac{r}{v}A_{-\eta+4} + \frac{1}{v}A_{-\eta+5},$$

for $\eta = 1, 2, 3, \dots$ where $v \neq 0$. Therefore, recurrence (1.1) satisfies for all integer η .

Over the last years, the Pentanacci sequence has been the subject of extensive research by various authors; see, for instance, [1], [2], [3], [4].

Table 1. Some special cases of the generalized Pentanacci sequences

No	Sequences (Numbers)	Notation	Ref
1	Generalized Pentanacci	$\{V_\eta\} = \{A_\eta(A_0, A_1, A_2, A_3, A_4; 1, 1, 1, 1, 1)\}$	[4]
2	Generalized Fifth order Pell	$\{V_\eta\} = \{A_\eta(A_0, A_1, A_2, A_3, A_4; 2, 1, 1, 1, 1)\}$	[5]
3	Generalized Fifth order Jacobsthal	$\{V_\eta\} = \{A_\eta(A_0, A_1, A_2, A_3, A_4; 1, 1, 1, 1, 2)\}$	[6]
4	Generalized 5-primes	$\{V_\eta\} = \{A_\eta(A_0, A_1, A_2, A_3, A_4; 2, 3, 5, 7, 11)\}$	[7]

For some specific values of A_0, A_1, A_2, A_3, A_4 and r, s, t, u, v , it is worthwhile to present these special Pentanacci numbers in a table under a specific name. As an example, the literature employs the sequence names and symbols (refer to Table 2) for specific cases of r, s, t, u, v , along with their initial values.

The sequences and notations used in this study are as follows: Pentanacci $\{P_\eta\}$, Pentanacci-Lucas $\{Q_\eta\}$, fifth-order Pell $\{P_\eta^{(5)}\}$, fifth-order Pell-Lucas $\{Q_\eta^{(5)}\}$, modified fifth-order Pell $\{E_\eta^{(5)}\}$, fifth-order Jacobsthal $\{J_\eta^{(5)}\}$, fifth-order Jacobsthal-Lucas $\{j_\eta^{(5)}\}$, modified fifth-order Jacobsthal $\{K_\eta^{(5)}\}$, fifth-order Jacobsthal Perrin $\{Q_\eta^{(5)}\}$, adjusted fifth-order Jacobsthal $\{S_\eta^{(5)}\}$, modified fifth-order Jacobsthal-Lucas $\{R_\eta^{(5)}\}$, 5-primes $\{G_\eta\}$, Lucas 5-primes $\{H_\eta\}$, modified 5-primes $\{E_\eta\}$.

To simplify notation, we henceforth omit the superscripts in these sequences. For instance, we use P_η instead of $P_\eta^{(5)}$.

Theorem 1.1. *Let z be a real (or complex) number. For $\eta \geq 0$, we have the following formulas:*

Table 2. Some members of generalized Pentanacci sequences

Notation	OEIS [8]	Ref
$\{P_\eta\} = \{A_\eta(0, 1, 1, 2, 4; 1, 1, 1, 1, 1)\}$	A001591	[4]
$\{Q_\eta\} = \{A_\eta(5, 1, 3, 7, 15; 1, 1, 1, 1, 1)\}$	A074048	[4]
$\{P_\eta^{(5)}\} = \{A_\eta(0, 1, 2, 5, 13; 2, 1, 1, 1, 1)\}$	A141448	[5]
$\{Q_\eta^{(5)}\} = \{A_\eta(5, 2, 6, 17, 46; 2, 1, 1, 1, 1)\}$		[5]
$\{E_\eta^{(5)}\} = \{A_\eta(0, 1, 1, 3, 8; 2, 1, 1, 1, 1)\}$		[5]
$\{J_\eta^{(5)}\} = \{A_\eta(0, 1, 1, 1, 1; 1, 1, 1, 1, 2)\}$	A226310	[6],[9]
$\{i_\eta^{(5)}\} = \{A_\eta(2, 1, 5, 10, 20; 1, 1, 1, 1, 2)\}$	A226311	[6],[9]
$\{K_\eta^{(5)}\} = \{A_\eta(3, 1, 3, 10, 20; 1, 1, 1, 1, 2)\}$		[6]
$\{Q_\eta^{(5)}\} = \{A_\eta(3, 0, 2, 8, 16; 1, 1, 1, 1, 2)\}$		[6]
$\{S_\eta^{(5)}\} = \{A_\eta(0, 1, 1, 2, 4; 1, 1, 1, 1, 2)\}$		[6]
$\{R_\eta^{(5)}\} = \{A_\eta(5, 1, 3, 7, 15; 1, 1, 1, 1, 2)\}$		[6]
$\{G_\eta\} = \{A_\eta(0, 0, 0, 1, 2; 2, 3, 5, 7, 11)\}$		[7]
$\{H_\eta\} = \{A_\eta(5, 2, 10, 41, 150; 2, 3, 5, 7, 11)\}$		[7]
$\{E_\eta\} = \{A_\eta(0, 0, 0, 1, 1; 2, 3, 5, 7, 11)\}$		[7]

Table 3. Some special studies of sum formulas

Sequences	Papers dealing with sum formulae
Pell and Pell-Lucas	[10],[11],[12],[13],[14]
Generalized Fibonacci	[15],[16],[17],[18],[19],[20],[21]
Generalized Tribonacci	[22],[23],[24]
Generalized Tetranacci	[25],[26],[27]
Generalized Pentanacci	[28],[29]
Generalized Hexanacci	[30],[31]

(a) If $rz + sz^2 + tz^3 + uz^4 + vz^5 - 1 \neq 0$, then

$$\sum_{\lambda=0}^{\eta} z^\lambda A_\lambda = \frac{\Theta_1(z)}{rz + sz^2 + tz^3 + uz^4 + vz^5 - 1} = \frac{\Theta_1(z)}{\Theta(z)},$$

where,

$$\Theta_1(z) = z^{\eta+4}A_{\eta+4} - (rz - 1)z^{\eta+3}A_{\eta+3} - (sz^2 + rz - 1)z^{\eta+2}A_{\eta+2} - (sz^2 + tz^3 + rz - 1)z^{\eta+1}A_{\eta+1} + vz^{\eta+5}A_\eta - z^4A_4 + z^3(rz - 1)A_3 + z^2(sz^2 + rz - 1)A_2 + z(sz^2 + tz^3 + rz - 1)A_1 + (sz^2 + tz^3 + uz^4 + rz - 1)A_0.$$

(b) If $r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1 \neq 0$, then

$$\sum_{\lambda=0}^{\eta} z^\lambda A_{2\lambda} = \frac{\Theta_2(z)}{r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1},$$

where,

$$\Theta_2(z) = -(uz^2 + sz - 1)z^{\eta+1}A_{2\eta+2} + (t + rs + vz + ruz)z^{\eta+2}A_{2\eta+1} + (u + t^2z - u^2z^2 + v^2z^3 + rt + 2tvz^2 + rvz - suz)z^{\eta+2}A_{2\eta} + (v + ru - suz + tuz)z^{\eta+2}A_{2\eta-1} + v(r + vz^2 + tz)z^{\eta+2}A_{2\eta-2} + z^2(uz^2 + sz - 1)A_4 - z^3(t + rs +$$

$$vz + ruz)A_3 + z(r^2z + uz^2 - s^2z^2 + 2sz + rtz^2 + rvz^3 - suz^3 - 1)A_2 - z^3(v + ru - svz + tuz)A_1 + (r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + 2sz + 2rtz^2 + rvz^3 - 2suz^3 + tvz^4 - 1)A_0.$$

(c) If $r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1 = 0$, then

$$\begin{aligned} & \sum_{\lambda=0}^{\eta} z^{\lambda} A_{2\lambda+1} \\ &= \frac{\Theta_3(z)}{r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1}, \end{aligned}$$

where,

$$\Theta_3(z) = (r + vz^2 + tz)z^{\eta+1}A_{2\eta+2} + (s - s^2z + t^2z^2 - u^2z^3 + v^2z^4 + uz + rvz^2 - 2suz^2 + 2tvz^3 + rtz)z^{\eta+1}A_{2\eta+1} + (t + vz - svz^2 + ruz - stz)z^{\eta+1}A_{2\eta} + (u - u^2z^2 + v^2z^3 + tvz^2 + rvz - suz)z^{\eta+1}A_{2\eta-1} - v(uz^2 + sz - 1)z^{\eta+1}A_{2\eta-2} - z^2(r + vz^2 + tz)A_4 + z(r^2z + uz^2 + sz + rtz^2 + rvz^3 - 1)A_3 - z^2(t + vz - svz^2 + ruz - stz)A_2 + (r^2z + uz^2 - s^2z^2 + t^2z^3 + 2sz + 2rtz^2 + rvz^3 - suz^3 + tvz^4 - 1)A_1 + vz^2(uz^2 + sz - 1)A_0$$

Proof. See Soykan [32], theorem 2.1, for the proof. □

Theorem 1.2. For $\eta \geq 1$, we have the following formulas: If $v + rz^4 + sz^3 + tz^2 + uz - z^5 \neq 0$, then

$$\sum_{\lambda=1}^{\eta} z^{\lambda} A_{-\lambda} = \frac{\Theta_4(z)}{v + rz^4 + sz^3 + tz^2 + uz - z^5},$$

where,

$$\Theta_4(z) = -z^{\eta+1}A_{4-\eta} + (r - z)z^{\eta+1}A_{-\eta+3} + (s + rz - z^2)z^{\eta+1}A_{-\eta+2} + (t + rz^2 + sz - z^3)z^{\eta+1}A_{-\eta+1} + (u + rz^3 + sz^2 + tz - z^4)z^{\eta+1}A_{-\eta} + zA_4 - z(r - z)A_3 + z(-s - rz + z^2)A_2 + z(-t - rz^2 - sz + z^3)A_1 + z(-u - rz^3 - sz^2 - tz + z^4)A_0.$$

Proof. See Soykan [32], theorem 4.1, for the proof. □

The paper is structured into 6 distinct sections. In section 1, we initially revisited the definition of generalized Pentanacci numbers, laying out the formal expressions and recurrence relations. We also drew from several important publications in the literature to build a foundation for our paper. For a better understanding, we present three detailed tables. They include notations, some members of sequences and related works. Moreover, we highlight two crucial theorems conducted by Soykan in [32], which have proven invaluable in this research. In Section 2, we compute the following sums for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with positive subscripts: $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$. In section 3, we provide the closed-form solutions (identities) for sums $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$ for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$. In section 4, we compute the following sum for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with negative subscripts: $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$. In section 5, we provide the closed-form solutions (identities) for sums $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$, for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$. Finally, section 6 is the conclusion. It summarises the findings of the article.

2 Sums $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$ with Non-negative Subscripts

During this section, we compute the following sums for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with positive subscripts: $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$. The following theorem presents three significant summation formulas for generalized Pentanacci numbers with positive subscripts.

Theorem 2.1. For $\eta \geq 0$ and let z be a real (or complex) number. Then, we get the following formulae:

(a) If $vz^5 + uz^4 + tz^3 + sz^2 + rz - 1 \neq 0$, then

$$\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda} = \frac{\Omega_1}{(vz^5 + uz^4 + tz^3 + sz^2 + rz - 1)^2},$$

where,

$$\begin{aligned} \Omega_1 = & z^{\eta+4}(\eta(sz^2 + tz^3 + uz^4 + vz^5 + rz - 1) - 4 + 2sz^2 + tz^3 - vz^5 + 3rz)A_{\eta+4} - z^{\eta+3}(\eta(rz - 1)(sz^2 + tz^3 + uz^4 + vz^5 + rz - 1) \\ & + 3 - sz^2 + uz^4 + 2vz^5 + 3r^2z^2 - 6rz + 2rsz^3 + rtz^4 - rvz^6)A_{\eta+3} - z^{\eta+2}(\eta(sz^2 + rz - 1)(sz^2 + tz^3 + uz^4 + vz^5 + rz - 1) \\ & + 2 - 4sz^2 + tz^3 + 2uz^4 + 3vz^5 + 2r^2z^2 + 2s^2z^4 - 4rz + 4rsz^3 - ruvz^5 + stz^5 - 2rvz^6 - svz^7)A_{\eta+2} - z^{\eta+1}(\eta(sz^2 + tz^3 + rz - 1)(sz^2 + tz^3 + uz^4 + vz^5 + rz - 1) \\ & + 1 + 2rsz^3 + 2rtz^4 - 2ruz^5 + 2stz^5 - 3rvz^6 - suz^6 - 2svz^7 - tvz^8 - 2sz^2 - 2tz^3 + 3uz^4 + 4vz^5 + r^2z^2 + s^2z^4 + t^2z^6 - 2rz)A_{\eta+1} + vz^{\eta+5}(\eta(sz^2 + tz^3 + uz^4 + vz^5 + rz - 1) + 3sz^2 + 2tz^3 + uz^4 + 4rz - 5) \\ & A_{\eta} - z^4(2sz^2 + tz^3 - vz^5 + 3rz - 4)A_4 + z^3(-sz^2 + uz^4 + 2vz^5 + 3r^2z^2 - 6rz + 2rsz^3 + rtz^4 - rvz^6 + 3) \\ & A_3 + z^2(-4sz^2 + tz^3 + 2uz^4 + 3vz^5 + 2r^2z^2 + 2s^2z^4 - 4rz + 4rsz^3 - ruvz^5 + stz^5 - 2rvz^6 - svz^7 + 2)A_2 + z(-2sz^2 - 2tz^3 + 3uz^4 + 4vz^5 + r^2z^2 + s^2z^4 + t^2z^6 - 2rz + 2rsz^3 + 2rtz^4 - 2ruz^5 + 2stz^5 - 3rvz^6 - suz^6 - 2svz^7 - tvz^8 + 1) \\ & A_1 - vz^5(3sz^2 + 2tz^3 + uz^4 + 4rz - 5)A_0. \end{aligned}$$

(b) If $r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1 \neq 0$, then

$$\begin{aligned} & \sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda} \\ & = \frac{\Omega_2}{(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1)^2}, \end{aligned}$$

where,

$$\begin{aligned} \Omega_2 = & -z^{\eta+1}(\eta(uz^2 + sz - 1)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) + 1 + s^2z^2 + 2t^2z^3 - u^2z^4 + u^3z^6 + 4v^2z^5 - 2sz + 2rtz^2 + 4rvz^3 + 6tvz^4 + 2r^2uz^3 - st^2z^4 + s^2uz^4 + 2su^2z^5 - 3sv^2z^6 - 2uv^2z^7 - 2rs \\ & vz^4 + 2rtuz^4 - 4stuvz^5 - 2tuvz^6 + r^2sz^2 - uz^2)A_{2\eta+2} + z^{\eta+2}(\eta(t + rs + vz + ruz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) + 2rs^2z - t^3z^3 - 2v^3z^6 - 2rs - 3vz - 2t + r^3sz + r^2tz + 4svz^2 + 2uvz^3 + 2ru^2z^3 + 2r^3uz^2 + 2r^2vz^2 + ru^3z^5 - s^2vz^3 + 2tu^2z^4 - 4t^2vz^4 - 5tv^2z^5 + u^2vz^5 + 2stuz^3 - rst^2z^3 + rs^2uz^3 + 2rsu^2z^4 - 2r^2svz^3 + 2r^2tuz^3 - 3rsv^2z^5 - 2ruv^2z^6 - 3ruz + 2stz + 4rsuz^2 - 4rstvz^4 - 2rtuvz^5)A_{2\eta+1} + z^{\eta+2}(\eta(u + t^2z - u^2z^2 + v^2z^3 + rt + 2tvz^2 + rvz - suz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) + 4u^2z^2 - 3t^2z - 2u - 2u^3z^4 - 5v^2z^3 - 2rt + 2r^2t^2z^2 - 3r^2u^2z^3 - s^2t^2z^3 + 4r^2v^2z^4 + 2s^2u^2z^4 - 3s^2v^2z^5 + r^3tz + r^2uz - 8tvz^2 + rt^3z^3 + 4st^2z^2 - 4s^2uz^2 - 6su^2z^3 + 2r^3vz^2 + s^3uz^3 + t^2uz^3 + su^3z^5 + 8sv^2z^4 + 2rv^3z^6 + 3uv^2z^5 + 4rsuv^2 + 12stvz^3 - 2r^2suz^2 - rs^2vz^3 - 2rtu^2z^4 + 6r^2tvz^3 + 4rt^2vz^4 + 5rtv^2z^5 - 4s^2tvz^4 - ru^2vz^5 - 2suw^2z^6 + 2rstz - 2stuvz^5 - 3rvz + 5suz + 4tuvz^4)A_{2\eta} + z^{\eta+2}(\eta(v + ru - svz + tuz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) + r^3uz - 3v^3z^5 - 2ru - 2v + r^2vz + 2ru^3z^4 - 2rv^2z^3 - 4s^2vz^2 + 2tu^2z^3 + s^3vz^3 - t^2vz^3 + tu^3z^5 - 4tv^2z^4 + 2sv^3z^6 + 2u^2vz^4 + 4stuz^2 + 2rsu^2z^3 - 2r^2svz^2 + 2r^2tuz^2 + rt^2uz^3 - s^2tuz^3 - 2r^2uvz^3 - 3ruv^2z^5 + 2stv^2z^5 - su^2vz^5 - 2t^2uvz^5 - 2tw^2z^6 - 2rstvz^3 - 4rtuvz^4 + 5svz - 3tuz + 2rsuz)A_{2\eta-1} + vz^{\eta+2}(\eta(r + vz^2 + tz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) + r^3z - 2r - 4vz^2 - v^3z^7 - 3tz + 4stz^2 + 6svz^3 + 2tuz^3 + 4uvz^4 + 2r^2tz^2 + rt^2z^3 - s^2tz^3 + 2ru^2z^4 + r^2vz^3 - rv^2z^5 - 2s^2vz^4 + tu^2z^5 - t^2vz^5 - 2tv^2z^6 + 2rsz + 2rsuz^3 - 2suvz^5)A_{2\eta-2} + z^2(-r^2z - 4uz^2 + 4s^2z^2 - s^3z^3 + t^2z^3 + 2u^2z^4 + 3v^2z^5 - 5sz + 2rvz^3 + 6suz^3 + 4tvz^4 + 2r^2sz^2 + 3r^2uz^3 - 2s^2uz^4 - su^2z^5 + t^2uz^5 - 2sv^2z^6 - uv^2z^7 + 2rstz^3 + 4rtuz^4 + 2rvuz^5 - 2stvz^5 + 2)A_4 + z^3(3t + v^3z^6 + 3rs + 4vz - 4rs^2z - 2r^3sz - 2r^2tz - 6svz^2 - 2tuz^2 - 4uvz^3 + rs^3z^2 - 2rt^2z^2 + s^2tz^2 - 4ru^2z^3 - 3r^3uz^2 - 3r^2vz^2 - 2rv^2z^4 + 2s^2vz^3 - tu^2z^4 + t^2vz^4 + 2tv^2z^5 + 4ruz - 4stz - 8rsuz^2 - 4rtvz^3 + 2suwz^4 - 2r^2stz^2 + 2rs^2uz^3 + rsu^2z^4 - 4r^2tuz^3 - rt^2uz^4 + 2rsv^2z^5 - 2r^2uvz^4 + ruv^2z^6 + 2rstvz^4)A_3 + z(r^4z^2 + 2r^3tz^3 + r^3vz^4 - 2r^2s^2z^3 - r^2su^2z^4 + 4r^2sz^2 + r^2t^2z^4 + 2r^2u^2z^5 + 2r^2uz^3 - r^2v^2z^6 - 2r^2z - 3rs^2z^4 - 2rs^2vz^5 - 4rstuz^5 + 4rstz^3 - 4rsuvz^6 + 2rsuvz^4 - rt^2vz^6 + rtu^2z^6 + 4rtuz^4 - 2rtv^2z^7 - rtz^2 + 4ruvz^5 - rv^3z^8 + s^4z^4 + 2s^3uz^5 - 4s^3z^3 + 2s^2tvz^6 + s^2u^2z^6 - 5s^2uz^4 + 2s^2v^2z^7 + 6s^2z^2 - st^2uz^6 - 2st^2z^4 - 8stvz^5 + suv^2z^8 + 4suz^3 - 6sv^2z^6 - 4sz + 2t^2z^3 - 2twz^6 + 6tvz^4 + u^3z^6 - u^2z^4 - 2uv^2z^7 - uz^2 + 4v^2z^5 + 1)A_2 + z^3(3v + 2v^3z^5 + 3ru - 2r^3uz - 2r^2vz - 2uvz^2 - 2ru^2z^2 - ru^3z^4 + \end{aligned}$$

$$7s^2vz^2 - 4tu^2z^3 - 2s^3vz^3 - t^3uz^4 + 2tv^2z^4 - sv^3z^6 - u^2vz^4 - 8svz + 4tuz - 2rtvz^2 - 6stuz^2 + 4suvz^3 + rs^2uz^2 + 3r^2svz^2 - 5r^2tuz^2 - 4rt^2uz^3 + 2rsv^2z^4 + 2s^2tuz^3 + 2stu^2z^4 + st^2vz^4 + 2ruv^2z^5 - 2s^2uvz^4 + tv^2z^6 - 4rsuz + 4rstvz^3) A_1 + vz^3(3r - 2r^3z + 5vz^2 - t^3z^4 + 4tz - 2ruz^2 - 6stz^2 - 8svz^3 - 4tuz^3 - 6uvz^4 + rs^2z^2 - 5r^2tz^2 - 4rt^2z^3 + 2s^2tz^3 - ru^2z^4 - 4r^2vz^3 - 2rv^2z^5 + 3s^2vz^4 - 2t^2vz^5 - tv^2z^6 + u^2vz^6 - 4rsz - 6rtvz^4 + 2stuz^4 + 4suvz^5)A_0.$$

(c) If $r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1 \neq 0$, then

$$\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1} = \frac{\Omega_3}{(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1)^2},$$

where,

$$\begin{aligned} \Omega_3 = & z^{\eta+1}(\eta(r + vz^2 + tz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) \\ & - 3vz^2 - t^3z^4 - 2v^3z^7 - 2tz - r - 2ruz^2 + 2stz^2 + 4svz^3 + 2uvz^4 + rs^2z^2 - r^2tz^2 - 2rt^2z^3 + 3ru^2z^4 - 2r^2vz^3 - 4rv^2z^5 - \\ & s^2vz^4 + 2tu^2z^5 - 4t^2vz^5 - 5tv^2z^6 + u^2vz^6 + 4rsuz^3 - 6rtvz^4 + 2stuz^4)A_{2\eta+2} + z^{\eta+1}(\eta(s - s^2z + t^2z^2 - u^2z^3 + v^2z^4 + \\ & uz + rvz^2 - 2suz^2 + 2tvz^3 + rtz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) \\ & + 2s^2z - s - s^3z^2 - 3t^2z^2 + 4u^2z^3 - 2u^3z^5 - 5v^2z^4 - r^2s^2z^2 + 2r^2t^2z^3 - 3r^2u^2z^4 + 4r^2v^2z^5 - 3rvz^2 + 6suz^2 - \\ & 8tvz^3 + r^3tz^2 + r^2uz^2 + rt^3z^4 + 2st^2z^3 - 4s^2uz^3 - 5su^2z^4 + 2r^3vz^3 + t^2uz^4 + 4sv^2z^5 + 2rv^3z^7 + 3uv^2z^6 + 6stvz^4 + \\ & 4tuvz^5 + rs^2vz^4 - 2rtu^2z^5 + 6r^2tvz^4 + 4rt^2vz^5 + 5rtv^2z^6 - ru^2vz^6 - 4r^2suz^3 - 2rstuz^4 - 2rtz - 2uz)A_{2\eta+1} + \\ & z^{\eta+1}(\eta(t + vz - svz^2 + ruz - stz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) \\ & + 5svz^2 - 2t^3z^3 - 3v^3z^6 - 2vz - t - 2tuz^2 - 2rt^2z^2 - s^2tz^2 + r^3uz^2 + r^2vz^2 + st^3z^4 + 2ru^3z^5 - 2rv^2z^4 - 4s^2vz^3 + \\ & 3tu^2z^4 + s^3vz^4 - 7t^2vz^4 - 8tv^2z^5 + 2sv^3z^7 + 2u^2vz^5 - 2ruz + 2stz + 2rsuz^2 - 4rtvz^3 + 4stuz^3 - r^2stz^2 - 2r^2svz^3 - rt^2 \\ & uz^4 - 2s^2tuz^4 - 2stu^2z^5 - 2r^2uvz^4 + 4st^2vz^5 - 3ruv^2z^6 + 5stv^2z^6 - su^2vz^6 + 2rsu^2z^4 - 4rtuvz^5)A_{2\eta} + z^{\eta+1}(\eta(u - \\ & u^2z^2 + v^2z^3 + tvz^2 + rvz - suz)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) \\ & + u^2z^2 - u + u^3z^4 - 4v^2z^3 - u^4z^6 - v^4z^8 - 2r^2u^2z^3 + r^2v^2z^4 - s^2u^2z^4 - 2s^2v^2z^5 - t^2v^2z^6 + 2u^2v^2z^7 - 3tvz^2 - s^2uz^2 + \\ & r^3vz^2 - 2t^2uz^3 - 2su^3z^5 + 6sv^2z^4 - rv^3z^6 - 2tv^3z^7 - 2rvz + 2suz + 2rsuz^2 - 2rtuz^2 - 4ruvz^3 + 4stvz^3 - 4tuvz^4 - r^2 \\ & suz^2 - 2rtu^2z^4 + 2r^2tvz^3 + rt^2vz^4 + st^2uz^4 - s^2tvz^4 + 2ru^2vz^5 + suv^2z^6 + 3tu^2vz^6 + 4rsuvz^4 + 4stuvz^5)A_{2\eta-1} - \\ & vz^{\eta+1}(\eta(uz^2 + sz - 1)(r^2z + 2uz^2 - s^2z^2 + t^2z^3 - u^2z^4 + v^2z^5 + 2sz + 2rtz^2 + 2rvz^3 - 2suz^3 + 2tvz^4 - 1) \\ & + 1 + s^2z^2 + 2t^2z^3 - uz^2 - u^2z^4 + u^3z^6 + 4v^2z^5 - 2sz + 2rtz^2 + 4rvz^3 + 6tvz^4 + r^2sz^2 + 2r^2uz^3 - st^2z^4 + \\ & s^2uz^4 + 2su^2z^5 - 3sv^2z^6 - 2uv^2z^7 - 2rsvz^4 + 2rtuz^4 - 4stvz^5 - 2tuvz^6)A_{2\eta-2} + z^2(2r - r^3z + 4vz^2 + v^3z^7 + 3tz - \\ & 4stz^2 - 6svz^3 - 2tuz^3 - 4uvz^4 - 2r^2tz^2 - rt^2z^3 + s^2tz^3 - 2ru^2z^4 - r^2vz^3 + rv^2z^5 + 2s^2vz^4 - tu^2z^5 + t^2vz^5 + 2tv^2z^6 - \\ & 2rsz - 2rsuz^3 + 2suvz^5)A_4 + z(r^4z^2 + 2r^3tz^3 + r^3vz^4 + 2r^2suz^4 + 3r^2sz^2 + r^2t^2z^4 + 2r^2u^2z^5 + 2r^2uz^3 - r^2v^2z^6 - \\ & 2r^2z - rs^2tz^4 - 2rs^2vz^5 + 4rstz^3 - 2rsuvz^6 + 4rsvz^4 - rt^2vz^6 + rtu^2z^6 + 4rtuz^4 - 2rtv^2z^7 - rtz^2 + 4ruvz^5 - \\ & rv^3z^8 + s^2uz^4 + s^2z^2 - st^2z^4 - 4stvz^5 + 2su^2z^5 - 3sv^2z^6 - 2sz + 2t^2z^3 - 2tuvz^6 + 6tvz^4 + u^3z^6 - u^2z^4 - \\ & 2uv^2z^7 - uz^2 + 4v^2z^5 + 1)A_3 + z^2(-2r^3uz^2 + 2r^2stz^2 + 3r^2svz^3 - 2r^2tuz^3 - r^2tz - 2r^2vz^2 + rs^2uz^3 + 2rst^2z^3 + 4rst \\ & vz^4 - 4rsuz^2 + 2rsuvz^5 + 2rtuvz^5 - ru^3z^5 - 2ru^2z^3 + 2ruv^2z^6 + 3ruz - s^3tz^3 - 2s^3vz^4 + 4s^2tz^2 - 2s^2uvz^5 + \\ & 7s^2vz^3 - st^2vz^5 + stu^2z^5 - 2stv^2z^6 - 5stz + 4suvz^4 - sv^3z^7 - 8svz^2 + t^3z^3 + 4t^2vz^4 - 2tu^2z^4 + 5tv^2z^5 + \\ & 2t - u^2vz^5 - 2uvz^3 + 2v^3z^6 + 3vz)A_2 + z^2(-2r^3vz^2 + 2r^2suz^2 - 5r^2tvz^3 + 3r^2u^2z^3 - r^2uz - 4r^2v^2z^4 + rs^2vz^3 + \\ & 2rstuz^3 - 4rsvz^2 - 4rt^2vz^4 + 4rtu^2z^4 - 6rtv^2z^5 + ru^2vz^5 - 2rv^3z^6 + 3rvz - s^3uz^3 + 2s^2tvz^4 - 2s^2u^2z^4 + 4s^2uz^2 + \\ & 3s^2v^2z^5 - 6stvz^3 - su^3z^5 + 6su^2z^3 + 2suv^2z^6 - 5suz - 8sv^2z^4 - t^3vz^5 + t^2u^2z^5 + t^2uz^3 - 2t^2v^2z^6 - tv^3z^7 + \\ & 4tvz^2 + 2u^3z^4 - 4u^2z^2 - 3uv^2z^5 + 2u + 5v^2z^3)A_1 + vz^2(-r^2z - 4uz^2 + 4s^2z^2 - s^3z^3 + t^2z^3 + 2u^2z^4 + 3v^2z^5 - 5sz + 2rvz^3 + \\ & 6suz^3 + 4tvz^4 + 2r^2sz^2 + 3r^2uz^3 - 2s^2uz^4 - su^2z^5 + t^2uz^5 - 2sv^2z^6 - uv^2z^7 + 2rstz^3 + 4rtuz^4 + 2ruvz^5 - 2stvz^5 + 2)A_0. \end{aligned}$$

Proof. (a) Using the following recurrence relation

$$A_{\eta} = rA_{\eta-1} + sA_{\eta-2} + tA_{\eta-3} + uA_{\eta-4} + vA_{\eta-5},$$

i.e.

$$vA_{\eta-5} = A_{\eta} - rA_{\eta-1} - sA_{\eta-2} - tA_{\eta-3} - uA_{\eta-4}.$$

We have

$$\begin{aligned}
 v \times 0 \times z^0 A_0 &= 0 \times z^0 A_5 - r \times 0 \times z^0 A_4 - s \times 0 \times z^0 A_3 - t \times 0 \times z^0 A_2 \\
 &\quad - u \times 0 \times z^0 A_1, \\
 v \times 1 \times z^1 A_1 &= 1 \times z^1 A_6 - r \times 1 \times z^1 A_5 - s \times 1 \times z^1 A_4 - t \times 1 \times z^1 A_3 \\
 &\quad - u \times 1 \times z^1 A_2, \\
 v \times 2 \times z^2 A_2 &= 2 \times z^2 A_7 - r \times 2 \times z^2 A_6 - s \times 2 \times z^2 A_5 - t \times 2 \times z^2 A_4 \\
 &\quad - u \times 2 \times z^2 A_3, \\
 &\quad \vdots \\
 v(\eta - 2)z^{\eta-2}A_{\eta-2} &= (\eta - 2)z^{\eta-2}A_{\eta+3} - r(\eta - 2)z^{\eta-2}A_{\eta+2} - s(\eta - 2)z^{\eta-2}A_{\eta+1} \\
 &\quad - t(\eta - 2)z^{\eta-2}A_{\eta} - u(\eta - 2)z^{\eta-2}A_{\eta-1}, \\
 v(\eta - 1)z^{\eta-1}A_{\eta-1} &= (\eta - 1)z^{\eta-1}A_{\eta+4} - r(\eta - 1)z^{\eta-1}A_{\eta+3} - s(\eta - 1)z^{\eta-1}A_{\eta+2} \\
 &\quad - t(\eta - 1)z^{\eta-1}A_{\eta+1} - u(\eta - 1)z^{\eta-1}A_{\eta}, \\
 v \times \eta \times z^{\eta}A_{\eta} &= \eta \times z^{\eta}A_{\eta+5} - r \times \eta \times z^{\eta}A_{\eta+4} - s \times \eta \times z^{\eta}A_{\eta+3} \\
 &\quad - t \times \eta \times z^{\eta}A_{\eta+2} - u \times \eta \times z^{\eta}A_{\eta+1}.
 \end{aligned}$$

By adding the equalities side by side and applying theorem 1.1 (a), we obtain (a).

(b) and (c) Using the following recurrence relation

$$A_{\eta} = rA_{\eta-1} + sA_{\eta-2} + tA_{\eta-3} + uA_{\eta-4} + vA_{\eta-5},$$

i.e.

$$rA_{\eta-1} = A_{\eta} - sA_{\eta-2} - tA_{\eta-3} - uA_{\eta-4} - vA_{\eta-5}.$$

We get

$$\begin{aligned}
 r \times 1 \times x^1 A_3 &= 1 \times x^1 A_4 - s \times 1 \times x^1 A_2 - t \times 1 \times x^1 A_1 \\
 &\quad - u \times 1 \times x^1 A_0 - v \times 1 \times x^1 A_{-1}, \\
 r \times 2 \times x^2 A_5 &= 2 \times x^2 A_6 - s \times 2 \times x^2 A_4 - t \times 2 \times x^2 A_3 \\
 &\quad - u \times 2 \times x^2 A_2 - v \times 2 \times x^2 A_1, \\
 &\quad \vdots \\
 r \times (\eta - 1) \times x^{\eta-1} A_{2\eta-1} &= (\eta - 1) \times x^{\eta-1} A_{2\eta} - s \times (\eta - 1) \times x^{\eta-1} A_{2\eta-2} \\
 &\quad - t \times (\eta - 1) \times x^{\eta-1} A_{2\eta-3} - u \times (\eta - 1) \times x^{\eta-1} A_{2\eta-4} \\
 &\quad - v \times (\eta - 1) \times x^{\eta-1} A_{2\eta-5}, \\
 r \times \eta \times x^{\eta} A_{2\eta+1} &= \eta \times x^{\eta} A_{2\eta+2} - s \times \eta \times x^{\eta} A_{2\eta} - t \times \eta \times x^{\eta} A_{2\eta-1} \\
 &\quad - u \times \eta \times x^{\eta} A_{2\eta-2} - v \times \eta \times x^{\eta} A_{2\eta-3}.
 \end{aligned}$$

By adding the equalities side by side the above equalities, we have that

$$\begin{aligned} r(-0 \times x^0 A_1 + \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda+1}) &= (\eta \times x^\eta A_{2\eta+2} - 0 \times x^0 A_2 - (-1) \times x^{-1} A_0 \\ &+ \sum_{\lambda=0}^{\eta} (\lambda - 1) x^{\lambda-1} A_{2\lambda}) - s(-0 \times x^0 A_0 + \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda}) - t(-(\eta + 1) x^{\eta+1} A_{2\eta+1} \\ &+ \sum_{\lambda=0}^{\eta} (\lambda + 1) x^{\lambda+1} A_{2\lambda+1}) - u(-(\eta + 1) x^{\eta+1} A_{2\eta} + \sum_{\lambda=0}^{\eta} (\lambda + 1) x^{\lambda+1} A_{2\lambda}) \\ &- v(-(\eta + 2) x^{\eta+2} A_{2\eta+1} - (\eta + 1) x^{\eta+1} A_{2\eta-1} + 1 \times x^1 A_{-1} \\ &+ \sum_{\lambda=0}^{\eta} (\lambda + 2) x^{\lambda+2} A_{2\lambda+1}). \end{aligned}$$

Since

$$A_{-1} = -\frac{u}{v} A_0 - \frac{t}{v} A_1 - \frac{s}{v} A_2 - \frac{r}{v} A_3 + \frac{1}{v} A_4,$$

we obtain

$$\begin{aligned} r(-0 \times x^0 A_1 + \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda+1}) &= (\eta \times x^\eta A_{2\eta+2} - 0 \times x^0 A_2 - (-1) \times x^{-1} A_0 \\ &+ x^{-1} \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda} - x^{-1} \sum_{\lambda=0}^{\eta} x^\lambda A_{2\lambda}) - s(-0 \times x^0 A_0 + \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda}) - t(-(\eta + 1) x^{\eta+1} A_{2\eta+1} \\ &+ x^1 \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda+1} + x^1 \sum_{\lambda=0}^{\eta} x^\lambda A_{2\lambda+1}) - u(-(\eta + 1) x^{\eta+1} A_{2\eta} + x^1 \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda} \\ &+ x^1 \sum_{\lambda=0}^{\eta} x^\lambda A_{2\lambda}) - v(-(\eta + 2) x^{\eta+2} A_{2\eta+1} - (\eta + 1) x^{\eta+1} A_{2\eta-1} + 1 \times x^1 (-\frac{u}{v} A_0 - \frac{t}{v} A_1 \\ &- \frac{s}{v} A_2 - \frac{r}{v} A_3 + \frac{1}{v} A_4) + x^2 \sum_{\lambda=0}^{\eta} \lambda x^\lambda A_{2\lambda+1} + 2x^2 \sum_{\lambda=0}^{\eta} x^\lambda A_{2\lambda+1}). \end{aligned} \tag{2.1}$$

In a similar way, using the following recurrence relation

$$A_\eta = rA_{\eta-1} + sA_{\eta-2} + tA_{\eta-3} + uA_{\eta-4} + vA_{\eta-5},$$

i.e.

$$rA_{\eta-1} = A_\eta - sA_{\eta-2} - tA_{\eta-3} - uA_{\eta-4} - vA_{\eta-5}.$$

We obtain the following obvious equalities;

$$\begin{aligned} r \times 1 \times x^1 A_2 &= 1 \times x^1 A_3 - s \times 1 \times x^1 A_1 - t \times 1 \times x^1 A_0 \\ &\quad - u \times 1 \times x^1 A_{-1} - v \times 1 \times x^1 A_{-2}, \\ r \times 2 \times x^2 A_4 &= 2 \times x^2 A_5 - s \times 2 \times x^2 A_3 - t \times 2 \times x^2 A_2 \\ &\quad - u \times 2 \times x^2 A_1 - v \times 2 \times x^2 A_0, \\ &\quad \vdots \\ r \times (\eta - 1) \times x^{\eta-1} A_{2\eta-2} &= (\eta - 1) \times x^{\eta-1} A_{2\eta-1} - s \times (\eta - 1) \times x^{\eta-1} A_{2\eta-3} \\ &\quad - t \times (\eta - 1) \times x^{\eta-1} A_{2\eta-4} - u \times (\eta - 1) \times x^{\eta-1} A_{2\eta-5} \\ &\quad - v \times (\eta - 1) \times x^{\eta-1} A_{2\eta-6}, \\ r \times \eta \times x^\eta A_{2\eta} &= \eta \times x^\eta A_{2\eta+1} - s \times \eta \times x^\eta A_{2\eta-1} \\ &\quad - t \times \eta \times x^\eta A_{2\eta-2} - u \times \eta \times x^\eta A_{2\eta-3} - v \times \eta \times x^\eta A_{2\eta-4}. \end{aligned}$$

By adding side by side the above equalities, we have that

$$\begin{aligned}
 r(-0 \times x^0 A_0 + \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda}) &= (-0 \times x^0 A_1 + \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda+1}) \\
 -s(-(\eta+1)x^{\eta+1} A_{2\eta+1} + \sum_{\lambda=0}^{\eta} (\lambda+1)x^{\lambda+1} A_{2\lambda+1}) &- t(-(\eta+1)x^{\eta+1} A_{2\eta}) \\
 + \sum_{\lambda=0}^{\eta} (\lambda+1)x^{\lambda+1} A_{2\lambda}) &- u(-(\eta+2)x^{\eta+2} A_{2\eta+1} - (\eta+1)x^{\eta+1} A_{2\eta-1}) \\
 + 1 \times x^1 A_{-1} + \sum_{\lambda=0}^{\eta} (\lambda+2)x^{\lambda+2} A_{2\lambda+1}) &- v(-(\eta+2)x^{\eta+2} A_{2\eta} - (\eta+1)x^{\eta+1} A_{2\eta-2}) \\
 + 1 \times x^1 A_{-2} + \sum_{\lambda=0}^{\eta} (\lambda+2)x^{\lambda+2} A_{2\lambda}). &
 \end{aligned}$$

Since

$$\begin{aligned}
 A_{-1} &= -\frac{u}{v} A_0 - \frac{t}{v} A_1 - \frac{s}{v} A_2 - \frac{r}{v} A_3 + \frac{1}{v} A_4, \\
 A_{-2} &= -\frac{u}{v} (-\frac{u}{v} A_0 - \frac{t}{v} A_1 - \frac{s}{v} A_2 - \frac{r}{v} A_3 + \frac{1}{v} A_4) - \frac{t}{v} A_0 - \frac{s}{v} A_1 - \frac{r}{v} A_2 + \frac{1}{v} A_3.
 \end{aligned}$$

We have that

$$\begin{aligned}
 r(-0 \times x^0 A_0 + \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda}) &= (-0 \times x^0 A_1 + \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda+1}) \tag{2.2} \\
 -s(-(\eta+1)x^{\eta+1} A_{2\eta+1} + x^1 \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda+1} + x^1 \sum_{\lambda=0}^{\eta} x^{\lambda} A_{2\lambda+1}) &- t(-(\eta+1)x^{\eta+1} A_{2\eta}) \\
 + x^1 \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda} + x^1 \sum_{\lambda=0}^{\eta} x^{\lambda} A_{2\lambda}) &- u(-(\eta+2)x^{\eta+2} A_{2\eta+1} - (\eta+1)x^{\eta+1} A_{2\eta-1}) \\
 + 1 \times x^1 (-\frac{u}{v} A_0 - \frac{t}{v} A_1 - \frac{s}{v} A_2 - \frac{r}{v} A_3 + \frac{1}{v} A_4) &+ x^2 \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda+1} + 2x^2 \sum_{\lambda=0}^{\eta} x^{\lambda} A_{2\lambda+1}) \\
 -v(-(\eta+2)x^{\eta+2} A_{2\eta} - (\eta+1)x^{\eta+1} A_{2\eta-2} + 1 \times x^1 (-\frac{u}{v} (-\frac{u}{v} A_0 - \frac{t}{v} A_1 - \frac{s}{v} A_2 - \frac{r}{v} A_3 & \\
 + \frac{1}{v} A_4) - \frac{t}{v} A_0 - \frac{s}{v} A_1 - \frac{r}{v} A_2 + \frac{1}{v} A_3) &+ x^2 \sum_{\lambda=0}^{\eta} \lambda x^{\lambda} A_{2\lambda} + 2x^2 \sum_{\lambda=0}^{\eta} x^{\lambda} A_{2\lambda}).
 \end{aligned}$$

Then, by using parts (b) and (c) of theorem 1.1, and solving the system of equations (2.1)-(2.2), the desired result follows. □

3 Results for Special z Values in Non-negative Subscripts

In this section, we handle the closed-form solutions (identities) for sums $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$ for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$.

3.1 The case $z = 1$

The case $z = 1$ of theorem 2.1 is provided in Soykan [33].

3.2 The case $z = -1$

During this subsection, we overcome the closed-form solutions (identities) for sums $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda+1}$ specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$.

Here, recall theorem 2.1 and set $r = s = t = u = v = 1$; this gives the result below.

Proposition 3.1. For $\eta \geq 0$ and setting $r, s, t, u, v = 1$ in theorem 2.1 leads to the next outcomes.

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{\lambda} = \frac{1}{4}((-1)^{\eta} (-(2\eta + 5)A_{\eta+4} + (4\eta + 8)A_{\eta+3} - (2\eta - 1)A_{\eta+2} + (4\eta + 2)A_{\eta+1} + (2\eta + 7)A_{\eta}) + 5A_4 - 8A_3 - A_2 - 2A_1 - 7A_0)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 1)A_{2\eta+2} + 4A_{2\eta+1} - (2\eta - 1)A_{2\eta} - (4\eta + 6)A_{2\eta-1} - (2\eta + 5)A_{2\eta-2}) + A_4 + 4A_3 - 5A_2 - 10A_1 - 7A_0)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda+1} = \frac{1}{4}((-1)^{\eta} ((2\eta + 3)A_{2\eta+2} - (2\eta + 7)A_{2\eta} - 6A_{2\eta-1} + (2\eta - 1)A_{2\eta-2}) + 5A_4 - 4A_3 - 9A_2 - 6A_1 + A_0)$.

The next corollary follows if we replace A_{η} in the proposition 3.1 with P_{η} . We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2$ and $P_4 = 4$.

Corollary 3.1. For $\eta \geq 0$, Pentanacci numbers get the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{\lambda} = \frac{1}{4}((-1)^{\eta} (-(2\eta + 5)P_{\eta+4} + (4\eta + 8)P_{\eta+3} - (2\eta - 1)P_{\eta+2} + (4\eta + 2)P_{\eta+1} + (2\eta + 7)P_{\eta}) + 1)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{2\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 1)P_{2\eta+2} + 4P_{2\eta+1} - (2\eta - 1)P_{2\eta} - (4\eta + 6)P_{2\eta-1} - (2\eta + 5)P_{2\eta-2}) - 3)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{2\lambda+1} = \frac{1}{4}((-1)^{\eta} ((2\eta + 3)P_{2\eta+2} - (2\eta + 7)P_{2\eta} - 6P_{2\eta-1} + (2\eta - 1)P_{2\eta-2}) - 3)$.

The next corollary follows if we replace A_{η} in the proposition 3.1 with Q_{η} . We additionally apply the initial conditions where $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7$ and $Q_4 = 15$.

Corollary 3.2. For $\eta \geq 0$, Pentanacci-Lucas numbers satisfy the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{\lambda} = \frac{1}{4}((-1)^{\eta} (-(2\eta + 5)Q_{\eta+4} + (4\eta + 8)Q_{\eta+3} - (2\eta - 1)Q_{\eta+2} + (4\eta + 2)Q_{\eta+1} + (2\eta + 7)Q_{\eta}) - 21)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{2\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 1)Q_{2\eta+2} + 4Q_{2\eta+1} - (2\eta - 1)Q_{2\eta} - (4\eta + 6)Q_{2\eta-1} - (2\eta + 5)Q_{2\eta-2}) - 17)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{2\lambda+1} = \frac{1}{4}((-1)^{\eta} ((2\eta + 3)Q_{2\eta+2} - (2\eta + 7)Q_{2\eta} - 6Q_{2\eta-1} + (2\eta - 1)Q_{2\eta-2}) + 19)$.

Here, recall theorem 2.1 and set $r = 2, s = t = u = v = 1$; this gives the result below.

Proposition 3.2. For $\eta \geq 0$ and setting $r = 2, s = t = u = v = 1$, in theorem 2.1 leads to the next outcomes.

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{\lambda} = \frac{1}{9}((-1)^{\eta} (-(3\eta + 8)A_{\eta+4} + (9\eta + 21)A_{\eta+3} - (6\eta + 4)A_{\eta+2} + (9\eta + 6)A_{\eta+1} + (3\eta + 11)A_{\eta}) + 8A_4 - 21A_3 + 4A_2 - 6A_1 - 11A_0)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda} = \frac{1}{25}((-1)^{\eta} ((5\eta - 6)A_{2\eta+2} + 15A_{2\eta+1} - (5\eta - 11)A_{2\eta} - (10\eta + 13)A_{2\eta-2} - (15\eta + 12)A_{2\eta-1}) - A_4 + 15A_3 - 19A_2 - 27A_1 - 23A_0)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda+1} = \frac{1}{25}((-1)^{\eta} ((10\eta + 3)A_{2\eta+2} + 5A_{2\eta+1} - (10\eta + 18)A_{2\eta} + (5\eta - 6)A_{2\eta-2} - (5\eta + 19)A_{2\eta-1}) + 13A_4 - 20A_3 - 28A_2 - 24A_1 - A_0)$.

The next corollary follows if we replace A_{η} in the proposition 3.2 with P_{η} . We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5$ and $P_4 = 13$.

Corollary 3.3. For $\eta \geq 0$, the fifth-order Pell numbers satisfy the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{\lambda} = \frac{1}{9}((-1)^{\eta}(-(3\eta+8)P_{\eta+4}+(9\eta+21)P_{\eta+3}-(6\eta+4)P_{\eta+2}+(9\eta+6)P_{\eta+1}+(3\eta+11)P_{\eta})+1)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{2\lambda} = \frac{1}{25}((-1)^{\eta}((5\eta-6)P_{2\eta+2}+15P_{2\eta+1}-(5\eta-11)P_{2\eta}-(10\eta+13)P_{2\eta-2}-(15\eta+12)P_{2\eta-1})-3)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} P_{2\lambda+1} = \frac{1}{25}((-1)^{\eta}((10\eta+3)P_{2\eta+2}+5P_{2\eta+1}-(10\eta+18)P_{2\eta}+(5\eta-6)P_{2\eta-2}-(5\eta+19)P_{2\eta-1})-11)$.

The next corollary follows if we replace A_{η} in the proposition 3.2 with P_{η} . We additionally apply the initial conditions where $Q_0 = 5, Q_1 = 2, Q_2 = 6, Q_3 = 17$ and $Q_4 = 46$.

Corollary 3.4. For $\eta \geq 0$, the fifth-order Pell-Lucas numbers have the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{\lambda} = \frac{1}{9}((-1)^{\eta}(-(3\eta+8)Q_{\eta+4}+(9\eta+21)Q_{\eta+3}-(6\eta+4)Q_{\eta+2}+(9\eta+6)Q_{\eta+1}+(3\eta+11)Q_{\eta})-32)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{2\lambda} = \frac{1}{25}((-1)^{\eta}((5\eta-6)Q_{2\eta+2}+15Q_{2\eta+1}-(5\eta-11)Q_{2\eta}-(10\eta+13)Q_{2\eta-2}-(15\eta+12)Q_{2\eta-1})-74)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} Q_{2\lambda+1} = \frac{1}{25}((-1)^{\eta}((10\eta+3)Q_{2\eta+2}+5Q_{2\eta+1}-(10\eta+18)Q_{2\eta}+(5\eta-6)Q_{2\eta-2}-(5\eta+19)Q_{2\eta-1})+37)$.

Here, recall theorem 2.1 and set $r = s = t = u = 1, v = 2$; this gives the result below.

Proposition 3.3. For $\eta \geq 0$ and setting $r, s, t, u = 1, v = 2$, in theorem 2.1 leads to the next outcomes.

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{\lambda} = \frac{1}{9}((-1)^{\eta}(-(3\eta+4)A_{\eta+4}+(6\eta+5)A_{\eta+3}-(3\eta-5)A_{\eta+2}+(6\eta-4)A_{\eta+1}+2(3\eta+7)A_{\eta})+4A_4-5A_3-5A_2+4A_1-14A_0)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda} = \frac{1}{25}((-1)^{\eta}((5\eta-14)A_{2\eta+2}+(5\eta+11)A_{2\eta+1}+(5\eta+36)A_{2\eta}-(20\eta-11)A_{2\eta-1}-2(10\eta+7)A_{2\eta-2})-9A_4+16A_3+16A_2-9A_1-34A_0)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda+1} = \frac{1}{25}((-1)^{\eta}((10\eta-3)A_{2\eta+2}+(10\eta+22)A_{2\eta+1}-(15\eta+3)A_{2\eta}+2(5\eta-14)A_{2\eta-2}-(15\eta+28)A_{2\eta-1})+7A_4+7A_3-18A_2-43A_1-18A_0)$.

The next corollary follows if we replace A_{η} in the proposition 3.3 with J_{η} . We additionally apply the initial conditions where $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1$ and $J_4 = 1$.

Corollary 3.5. For $\eta \geq 0$, the fifth-order Jacobsthal numbers get the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} J_{\lambda} = \frac{1}{9}((-1)^{\eta}(-(3\eta+4)J_{\eta+4}+(6\eta+5)J_{\eta+3}-(3\eta-5)J_{\eta+2}+(6\eta-4)J_{\eta+1}+2(3\eta+7)J_{\eta})-2)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} J_{2\lambda} = \frac{1}{25}((-1)^{\eta}((5\eta-14)J_{2\eta+2}+(5\eta+11)J_{2\eta+1}+(5\eta+36)J_{2\eta}-(20\eta-11)J_{2\eta-1}-2(10\eta+7)J_{2\eta-2})+14)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} J_{2\lambda+1} = \frac{1}{25}((-1)^{\eta}((10\eta-3)J_{2\eta+2}+(10\eta+22)J_{2\eta+1}-(15\eta+3)J_{2\eta}+2(5\eta-14)J_{2\eta-2}-(15\eta+28)J_{2\eta-1})-47)$.

The next corollary follows if we replace A_{η} in the proposition 3.3 with j_{η} . We additionally apply the initial conditions where $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10$ and $j_4 = 20$.

Corollary 3.6. For $\eta \geq 0$, the fifth-order Jacobsthal-Lucas numbers satisfy the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} j_{\lambda} = \frac{1}{9}((-1)^{\eta}(-(3\eta+4)j_{\eta+4}+(6\eta+5)j_{\eta+3}-(3\eta-5)j_{\eta+2}+(6\eta-4)j_{\eta+1}+2(3\eta+7)j_{\eta})-19)$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} j_{2\lambda} = \frac{1}{25}((-1)^{\eta}((5\eta-14)j_{2\eta+2}+(5\eta+11)j_{2\eta+1}+(5\eta+36)j_{2\eta}-(20\eta-11)j_{2\eta-1}-2(10\eta+7)j_{2\eta-2})-17)$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} j_{2\lambda+1} = \frac{1}{25}((-1)^{\eta}((10\eta-3)j_{2\eta+2}+(10\eta+22)j_{2\eta+1}-(15\eta+3)j_{2\eta}+2(5\eta-14)j_{2\eta-2}-(15\eta+28)j_{2\eta-1})+41)$.

The next corollary follows if we replace A_η in the proposition 3.3 with K_η . We additionally apply the initial conditions where $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10$ and $K_4 = 20$.

Corollary 3.7. For $\eta \geq 0$, the modified fifth-order Jacobsthal numbers have the following property:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda K_\lambda = \frac{1}{9}((-1)^\eta (-3\eta+4)K_{\eta+4} + (6\eta+5)K_{\eta+3} - (3\eta-5)K_{\eta+2} + (6\eta-4)K_{\eta+1} + 2(3\eta+7)K_\eta) - 23$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda K_{2\lambda} = \frac{1}{25}((-1)^\eta ((5\eta-14)K_{2\eta+2} + (5\eta+11)K_{2\eta+1} + (5\eta+36)K_{2\eta} - (20\eta-11)K_{2\eta-1} - 2(10\eta+7)K_{2\eta-2}) - 83$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda K_{2\lambda+1} = \frac{1}{25}((-1)^\eta ((10\eta-3)K_{2\eta+2} + (10\eta+22)K_{2\eta+1} - (15\eta+3)K_{2\eta} + 2(5\eta-14)K_{2\eta-2} - (15\eta+28)K_{2\eta-1}) + 59$.

The next corollary follows if we replace A_η in the proposition 3.3 with Q_η . We additionally apply the initial conditions where $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8$ and $Q_4 = 16$.

Corollary 3.8. For $\eta \geq 0$, the fifth-order Jacobsthal Perrin numbers get the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda Q_\lambda = \frac{1}{9}((-1)^\eta (-3\eta+4)Q_{\eta+4} + (6\eta+5)Q_{\eta+3} - (3\eta-5)Q_{\eta+2} + (6\eta-4)Q_{\eta+1} + 2(3\eta+7)Q_\eta) - 28$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda Q_{2\lambda} = \frac{1}{25}((-1)^\eta ((5\eta-14)Q_{2\eta+2} + (5\eta+11)Q_{2\eta+1} + (5\eta+36)Q_{2\eta} - (20\eta-11)Q_{2\eta-1} - 2(10\eta+7)Q_{2\eta-2}) - 86$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda Q_{2\lambda+1} = \frac{1}{25}((-1)^\eta ((10\eta-3)Q_{2\eta+2} + (10\eta+22)Q_{2\eta+1} - (15\eta+3)Q_{2\eta} + 2(5\eta-14)Q_{2\eta-2} - (15\eta+28)Q_{2\eta-1}) + 78$.

The next corollary follows if we replace A_η in the proposition 3.3 with S_η . We additionally apply the initial conditions where $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2$ and $S_4 = 4$.

Corollary 3.9. For $\eta \geq 0$, the adjusted fifth-order Jacobsthal numbers satisfy the followings:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda S_\lambda = \frac{1}{9}((-1)^\eta (-3\eta+4)S_{\eta+4} + (6\eta+5)S_{\eta+3} - (3\eta-5)S_{\eta+2} + (6\eta-4)S_{\eta+1} + 2(3\eta+7)S_\eta) + 5$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda S_{2\lambda} = \frac{1}{25}((-1)^\eta ((5\eta-14)S_{2\eta+2} + (5\eta+11)S_{2\eta+1} + (5\eta+36)S_{2\eta} - (20\eta-11)S_{2\eta-1} - 2(10\eta+7)S_{2\eta-2}) + 3$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda S_{2\lambda+1} = \frac{1}{25}((-1)^\eta ((10\eta-3)S_{2\eta+2} + (10\eta+22)S_{2\eta+1} - (15\eta+3)S_{2\eta} + 2(5\eta-14)S_{2\eta-2} - (15\eta+28)S_{2\eta-1}) - 19$.

The next corollary follows if we replace A_η in the proposition 3.3 with R_η . We additionally apply the initial conditions where $R_0 = 5, R_1 = 1, R_2 = 3, R_3 = 7$ and $R_4 = 15$.

Corollary 3.10. The following properties is satisfied by the modified fifth-order Jacobsthal-Lucas numbers, for $\eta \geq 0$:

- (a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda R_\lambda = \frac{1}{9}((-1)^\eta (-3\eta+4)R_{\eta+4} + (6\eta+5)R_{\eta+3} - (3\eta-5)R_{\eta+2} + (6\eta-4)R_{\eta+1} + 2(3\eta+7)R_\eta) - 56$.
- (b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda R_{2\lambda} = \frac{1}{25}((-1)^\eta ((5\eta-14)R_{2\eta+2} + (5\eta+11)R_{2\eta+1} + (5\eta+36)R_{2\eta} - (20\eta-11)R_{2\eta-1} - 2(10\eta+7)R_{2\eta-2}) - 154$.
- (c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^\lambda R_{2\lambda+1} = \frac{1}{25}((-1)^\eta ((10\eta-3)R_{2\eta+2} + (10\eta+22)R_{2\eta+1} - (15\eta+3)R_{2\eta} + 2(5\eta-14)R_{2\eta-2} - (15\eta+28)R_{2\eta-1}) - 33$.

Here, recall theorem 2.1 and set $r = 2, s = 3, t = 5, u = 7, v = 11$; this gives the result below.

Proposition 3.4. For $\eta \geq 0$ and setting $r = 2, s = 3, t = 5, u = 7, v = 11$, in theorem 2.1 leads to the next outcomes.

(a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{\lambda} = \frac{1}{81}((-1)^{\eta}(-9\eta-2)A_{\eta+4} + (27\eta-15)A_{\eta+3} + 36A_{\eta+2} + (45\eta-46)A_{\eta+1} + 11(9\eta+7)A_{\eta}) - 2A_4 + 15A_3 - 36A_2 + 46A_1 - 77A_0$.

(b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda} = \frac{1}{5329}((-1)^{\eta}(-219\eta-184)A_{2\eta+2} + (1022\eta-1759)A_{2\eta+1} + (5037\eta+1827)A_{2\eta} - (1679\eta-6034)A_{2\eta-1} - 11(584\eta-807)A_{2\eta-2}) - 35A_4 - 737A_3 + 1535A_2 + 4355A_1 + 2453A_0$.

(c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} A_{2\lambda+1} = \frac{1}{5329}((-1)^{\eta}((584\eta-1391)A_{2\eta+2} + (4380\eta+2379)A_{2\eta+1} - (2774\eta-6954)A_{2\eta} - (7957\eta-10165)A_{2\eta-1} - 11(219\eta-184)A_{2\eta-2}) - 807A_4 + 1430A_3 + 4180A_2 + 2208A_1 - 385A_0$.

The next corollary follows if we replace A_{η} in the proposition 3.4 with G_{η} . We additionally apply the initial conditions where $G_0 = 0, G_1 = 0, G_2 = 0, G_3 = 1$ and $G_4 = 2$.

Corollary 3.11. For $\eta \geq 0$, the 5-primes numbers satisfy the followings:

(a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} G_{\lambda} = \frac{1}{81}((-1)^{\eta}(-9\eta-2)G_{\eta+4} + (27\eta-15)G_{\eta+3} + 36G_{\eta+2} + (45\eta-46)G_{\eta+1} + 11(9\eta+7)G_{\eta} + 11)$.

(b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} G_{2\lambda} = \frac{1}{5329}((-1)^{\eta}(-219\eta-184)G_{2\eta+2} + (1022\eta-1759)G_{2\eta+1} + (5037\eta+1827)G_{2\eta} - (1679\eta-6034)G_{2\eta-1} - 11(584\eta-807)G_{2\eta-2}) - 807$.

(c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} G_{2\lambda+1} = \frac{1}{5329}((-1)^{\eta}((584\eta-1391)G_{2\eta+2} + (4380\eta+2379)G_{2\eta+1} - (2774\eta-6954)G_{2\eta} - (7957\eta-10165)G_{2\eta-1} - 11(219\eta-184)G_{2\eta-2}) - 184$.

The next corollary follows if we replace A_{η} in the proposition 3.4 with H_{η} . We additionally apply the initial conditions where $H_0 = 5, H_1 = 2, H_2 = 10, H_3 = 41$ and $H_4 = 150$.

Corollary 3.12. The following properties is satisfied by the Lucas 5-primes numbers, for $\eta \geq 0$:

(a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} H_{\lambda} = \frac{1}{81}((-1)^{\eta}(-9\eta-2)H_{\eta+4} + (27\eta-15)H_{\eta+3} + 36H_{\eta+2} + (45\eta-46)H_{\eta+1} + 11(9\eta+7)H_{\eta}) - 338$.

(b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} H_{2\lambda} = \frac{1}{5329}((-1)^{\eta}(-219\eta-184)H_{2\eta+2} + (1022\eta-1759)H_{2\eta+1} + (5037\eta+1827)H_{2\eta} - (1679\eta-6034)H_{2\eta-1} - 11(584\eta-807)H_{2\eta-2}) + 858$.

(c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} H_{2\lambda+1} = \frac{1}{5329}((-1)^{\eta}((584\eta-1391)H_{2\eta+2} + (4380\eta+2379)H_{2\eta+1} - (2774\eta-6954)H_{2\eta} - (7957\eta-10165)H_{2\eta-1} - 11(219\eta-184)H_{2\eta-2}) - 18129$.

The next corollary follows if we replace A_{η} in the proposition 3.4 with E_{η} . We additionally apply the initial conditions where $E_0 = 0, E_1 = 0, E_2 = 0, E_3 = 1$ and $E_4 = 1$.

Corollary 3.13. For $\eta \geq 0$, the modified 5-primes numbers get the followings:

(a) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} E_{\lambda} = \frac{1}{81}((-1)^{\eta}(-9\eta-2)E_{\eta+4} + (27\eta-15)E_{\eta+3} + 36E_{\eta+2} + (45\eta-46)E_{\eta+1} + 11(9\eta+7)E_{\eta}) + 13$.

(b) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} E_{2\lambda} = \frac{1}{5329}((-1)^{\eta}(-219\eta-184)E_{2\eta+2} + (1022\eta-1759)E_{2\eta+1} + (5037\eta+1827)E_{2\eta} - (1679\eta-6034)E_{2\eta-1} - 11(584\eta-807)E_{2\eta-2}) - 772$.

(c) $\sum_{\lambda=0}^{\eta} \lambda(-1)^{\lambda} E_{2\lambda+1} = \frac{1}{5329}((-1)^{\eta}((584\eta-1391)E_{2\eta+2} + (4380\eta+2379)E_{2\eta+1} - (2774\eta-6954)E_{2\eta} - (7957\eta-10165)E_{2\eta-1} - 11(219\eta-184)E_{2\eta-2}) + 623$.

3.3 The case $z = i$

During this subsection, we cope with the closed-form solutions (identities) for sums $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{\lambda}$, $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{2\lambda+1}$ specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$. Here, recall theorem 2.1 and set $z = i$, $r = s = t = u = v = 1$; this gives the result below.

Proposition 3.5. For $\eta \geq 0$ and setting $r, s, t, u, v = 1$, in theorem 2.1 leads to the next outcomes.

(a) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{\lambda} = \frac{1}{2i}(i^{\eta}(((1-i)\eta + 6 - i)A_{\eta+4} - 2(\eta + (3+2i))A_{\eta+3} + (5i - 10 - (1-3i)\eta)A_{\eta+2} + ((2+2i)\eta - 4 + 10i)A_{\eta+1} + (((1+i)\eta + 2 + 7i))A_{\eta}) - (6-i)A_4 + (6+4i)A_3 + (10-5i)A_2 + (4-10i)A_1 - (2+7i)A_0).$

(b) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{2\lambda} = \frac{1}{18i}(i^{\eta}(((9+3i)\eta + 6 + 7i)A_{2\eta+2} - (12\eta + 16 + 12i)A_{2\eta+1} + (8+i - (3-3i)\eta)A_{2\eta} + (2i - 8 - (6-6i)\eta)A_{2\eta-1} + (2-13i - (3+3i)\eta)A_{2\eta-2}) + (10-15i)A_4 - (12-28i)A_3 - (14+5i)A_2 + (8+14i)A_1 - (16-i)A_0).$

(c) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} A_{2\lambda+1} = \frac{1}{18i}(i^{\eta}(((3i-3)\eta - 10 - 5i)A_{2\eta+2} + ((6+6i)\eta + 14 + 8i)A_{2\eta+1} + ((3+9i)\eta - 2 + 9i)A_{2\eta} + (6\eta + 8 - 6i)A_{2\eta-1} + ((9+3i)\eta + 6 + 7i)A_{2\eta-2}) - (2-13i)A_4 - (4+20i)A_3 + (18-i)A_2 - (6+14i)A_1 + (10-15i)A_0).$

The next corollary follows if we replace A_{η} in the proposition 3.5 with P_{η} . We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2$ and $P_4 = 4$.

Corollary 3.14. For $\eta \geq 0$, the Pentanacci numbers satisfy the followings.

(a) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} P_{\lambda} = \frac{1}{2i}(i^{\eta}(((1-i)\eta + 6 - i)P_{\eta+4} - 2(\eta + (3+2i))P_{\eta+3} + (5i - 10 - (1-3i)\eta)P_{\eta+2} + ((2+2i)\eta - 4 + 10i)P_{\eta+1} + (((1+i)\eta + 2 + 7i))P_{\eta}) + (2-3i)).$

(b) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} P_{2\lambda} = \frac{1}{18i}(i^{\eta}(((9+3i)\eta + 6 + 7i)P_{2\eta+2} - (12\eta + 16 + 12i)P_{2\eta+1} + (8+i - (3-3i)\eta)P_{2\eta} + (2i - 8 - (6-6i)\eta)P_{2\eta-1} + (2-13i - (3+3i)\eta)P_{2\eta-2}) + (10+5i)).$

(c) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} P_{2\lambda+1} = \frac{1}{18i}(i^{\eta}(((3i-3)\eta - 10 - 5i)P_{2\eta+2} + ((6+6i)\eta + 14 + 8i)P_{2\eta+1} + ((3+9i)\eta - 2 + 9i)P_{2\eta} + (6\eta + 8 - 6i)P_{2\eta-1} + ((9+3i)\eta + 6 + 7i)P_{2\eta-2}) + (-4-3i)).$

The next corollary follows if we replace A_{η} in the proposition 3.5 with Q_{η} . We additionally apply the initial conditions where $Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7$ and $Q_4 = 15$.

Corollary 3.15. For $\eta \geq 0$, the Pentanacci-Lucas numbers get the followings.

(a) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} Q_{\lambda} = \frac{1}{2i}(i^{\eta}(((1-i)\eta + 6 - i)Q_{\eta+4} - 2(\eta + (3+2i))Q_{\eta+3} + (5i - 10 - (1-3i)\eta)Q_{\eta+2} + ((2+2i)\eta - 4 + 10i)Q_{\eta+1} + (((1+i)\eta + 2 + 7i))Q_{\eta}) + (-24 - 17i)).$

(b) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} Q_{2\lambda} = \frac{1}{18i}(i^{\eta}(((9+3i)\eta + 6 + 7i)Q_{2\eta+2} - (12\eta + 16 + 12i)Q_{2\eta+1} + (8+i - (3-3i)\eta)Q_{2\eta} + (2i - 8 - (6-6i)\eta)Q_{2\eta-1} + (2-13i - (3+3i)\eta)Q_{2\eta-2}) + (-48 - 25i)).$

(c) $\sum_{\lambda=0}^{\eta} \lambda i^{\lambda} Q_{2\lambda+1} = \frac{1}{18i}(i^{\eta}(((3i-3)\eta - 10 - 5i)Q_{2\eta+2} + ((6+6i)\eta + 14 + 8i)Q_{2\eta+1} + ((3+9i)\eta - 2 + 9i)Q_{2\eta} + (6\eta + 8 - 6i)Q_{2\eta-1} + ((9+3i)\eta + 6 + 7i)Q_{2\eta-2}) + (40 - 37i)).$

In a similar way, readers can calculate the corresponding sums of other generalized fifth-order Pentanacci numbers.

4 Sum $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$ with Negative Subscripts

In this section, we compute the following sum for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with negative subscripts: $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$.

Theorem 4.1. For $\eta \geq 1$ and let z be a real (or complex) number. Then, we get the following formula: If $v + rz^4 + sz^3 + tz^2 + uz - z^5 \neq 0$, then

$$\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda} = \frac{\Omega_4}{(v + rz^4 + sz^3 + tz^2 + uz - z^5)^2},$$

where,

$$\begin{aligned} \Omega_4 = & z^{\eta+1}(\eta(-v-rz^4-sz^3-tz^2-uz+z^5)-v+3rz^4+2sz^3+tz^2-4z^5)A_{-\eta+4}+z^{\eta+1}(\eta(r-z)(v+rz^4+sz^3+tz^2+uz-z^5)+6rz^5+sz^4-uz^2-3r^2z^4+rv-2vz-3z^6-2rsz^3-rtz^2)A_{-\eta+3}+z^{\eta+1}(\eta(s+rz-z^2)(v+rz^4+sz^3+tz^2+uz-z^5)+4rz^6+4sz^5-tz^4-2uz^3-3vz^2-2r^2z^5-2s^2z^3+sv-2z^7-4rsz^4+ruz^2-stz^2+2rvz)A_{-\eta+2}+z^{\eta+1}(\eta(t+rz^2+sz-z^3)(v+rz^4+sz^3+tz^2+uz-z^5)+2rz^7+2sz^6+2tz^5-3uz^4-4vz^3-r^2z^6-s^2z^4-t^2z^2+tv-z^8-2rsz^5-2rtz^4+2ruz^3-2stz^3+3rvz^2+su^2+2svz)A_{-\eta+1}+z^{\eta+1}(\eta(u+rz^3+sz^2+tz-z^4)(v+rz^4+sz^3+tz^2+uz-z^5)-5vz^4+uv+4rvz^3+3svz^2+2tvz)A_{-\eta}+z(v-3rz^4-2sz^3-tz^2+4z^5)A_4+z(-6rz^5-sz^4+uz^2+3r^2z^4-rv+2vz+3z^6+2rsz^3+rtz^2)A_3+z(-4rz^6-4sz^5+tz^4+2uz^3+3vz^2+2r^2z^5+2s^2z^3-sv+2z^7+4rsz^4-ruz^2+stz^2-2rvz)A_2+z(-2rz^7-2sz^6-2tz^5+3uz^4+4vz^3+r^2z^6+s^2z^4+t^2z^2-tv+z^8+2rsz^5+2rtz^4-2ruz^3+2stz^3-3rvz^2-suz^2-2svz)A_1+vz(-u-4rz^3-3sz^2-2tz+5z^4)A_0. \end{aligned}$$

Proof. Using the following recurrence relation

$$\begin{aligned} A_{\eta+5} &= rA_{\eta+4} + sA_{\eta+3} + tA_{\eta+2} + uA_{\eta+1} + vA_{\eta} \\ \Rightarrow A_{-\eta+5} &= rA_{-\eta+4} + sA_{-\eta+3} + tA_{-\eta+2} + uA_{-\eta+1} + vA_{-\eta} \\ \Rightarrow A_{-\eta} &= -\frac{u}{v}A_{-\eta+1} - \frac{t}{v}A_{-\eta+2} - \frac{s}{v}A_{-\eta+3} - \frac{r}{v}A_{-\eta+4} + \frac{1}{v}A_{-\eta+5} \\ \Rightarrow A_{-\eta} &= \frac{1}{v}A_{-\eta+5} - \frac{u}{v}A_{-\eta+1} - \frac{t}{v}A_{-\eta+2} - \frac{s}{v}A_{-\eta+3} - \frac{r}{v}A_{-\eta+4} \end{aligned}$$

i.e.

$$vA_{-\eta} = A_{-\eta+5} - rA_{-\eta+4} - sA_{-\eta+3} - tA_{-\eta+2} - uA_{-\eta+1}.$$

We obtain

$$\begin{aligned} v \times \eta \times z^{\eta} A_{-\eta} &= \eta \times z^{\eta} A_{-\eta+5} - r \times \eta \times z^{\eta} A_{-\eta+4} \\ &\quad - s \times \eta \times z^{\eta} A_{-\eta+3} - t \times \eta \times z^{\eta} A_{-\eta+2} - u \times \eta \times z^{\eta} A_{-\eta+1}, \\ v(\eta-1)z^{\eta-1}A_{-\eta+1} &= (\eta-1)z^{\eta-1}A_{-\eta+6} - r(\eta-1)z^{\eta-1}A_{-\eta+5} \\ &\quad - s(\eta-1)z^{\eta-1}A_{-\eta+4} - t(\eta-1)z^{\eta-1}A_{-\eta+3} - u(\eta-1)z^{\eta-1}A_{-\eta+2}, \\ v(\eta-2)z^{\eta-2}A_{-\eta+2} &= (\eta-2)z^{\eta-2}A_{-\eta+7} - r(\eta-2)z^{\eta-2}A_{-\eta+6} \\ &\quad - s(\eta-2)z^{\eta-2}A_{-\eta+5} - t(\eta-2)z^{\eta-2}A_{-\eta+4} - u(\eta-2)z^{\eta-2}A_{-\eta+3}, \\ &\quad \vdots \\ v \times 3 \times z^3 A_{-3} &= 3 \times z^3 A_2 - r \times 3 \times z^3 A_1 \\ &\quad - s \times 3 \times z^3 A_0 - t \times 3 \times z^3 A_{-1} - u \times 3 \times z^3 A_{-2}, \\ v \times 2 \times z^2 A_{-2} &= 2 \times z^2 A_3 - r \times 2 \times z^2 A_2 \\ &\quad - s \times 2 \times z^2 A_1 - t \times 2 \times z^2 A_0 - u \times 2 \times z^2 A_{-1}, \\ v \times 1 \times z^1 A_{-1} &= 1 \times z^1 A_4 - r \times 1 \times z^1 A_3 \\ &\quad - s \times 1 \times z^1 A_2 - t \times 1 \times z^1 A_1 - u \times 1 \times z^1 A_0. \end{aligned}$$

By adding the identities side by side and applying theorem 1.2 (a), we arrive the result. □

5 Results for Special z Values in Negative Subscripts

This section provides the closed-form solutions (identities) for sums $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$, for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$.

5.1 The case $z = 1$

See Soykan [33] for the case $z = 1$ of theorem 4.1.

5.2 The case $z = -1$

This subsection considers the special case $z = -1$.

In this subsection, we give the closed-form solutions (identities) for sums $\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} A_{-\lambda}$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$.

Now, recall theorem 4.1 and set $r = s = t = u = v = 1$; this gives the result below.

Proposition 5.1. For $\eta \geq 1$ and setting $r, s, t, u, v = 1$, in theorem 4.1 leads to the next outcome.

$$\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} A_{-\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 5)A_{-\eta+4} - (4\eta - 8)A_{-\eta+3} + (2\eta + 1)A_{-\eta+2} - (4\eta - 2)A_{-\eta+1} + (2\eta + 7)A_{-\eta}) + 5A_4 - 8A_3 - A_2 - 2A_1 - 7A_0).$$

The next corollary follows if we replace A_{η} in the proposition 5.1 with P_{η} and Q_{η} respectively. We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 1, P_3 = 2, P_4 = 4, Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7$ and $Q_4 = 15$.

Corollary 5.1. For $\eta \geq 1$, we get the followings.

(a) $\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} P_{-\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 5)P_{-\eta+4} - (4\eta - 8)P_{-\eta+3} + (2\eta + 1)P_{-\eta+2} - (4\eta - 2)P_{-\eta+1} + (2\eta + 7)P_{-\eta}) + 1).$

(b) $\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} Q_{-\lambda} = \frac{1}{4}((-1)^{\eta} ((2\eta - 5)Q_{-\eta+4} - (4\eta - 8)Q_{-\eta+3} + (2\eta + 1)Q_{-\eta+2} - (4\eta - 2)Q_{-\eta+1} + (2\eta + 7)Q_{-\eta}) - 21).$

Here, recall theorem 4.1 and set $r = 2, s = t = u = v = 1$; this gives the result below.

Proposition 5.2. For $\eta \geq 1$ and setting $r = 2$ and $s, t, u, v = 1$, in theorem 4.1 leads to the next outcome.

$$\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} A_{-\lambda} = \frac{1}{9}((-1)^{\eta} ((3\eta - 8)A_{-\eta+4} - (9\eta - 21)A_{-\eta+3} + (6\eta - 4)A_{-\eta+2} - (9\eta - 6)A_{-\eta+1} + (6\eta + 11)A_{-\eta}) + 8A_4 - 21A_3 + 4A_2 - 6A_1 - 11A_0).$$

The next corollary follows if we replace A_{η} in the proposition 5.2 with P_{η} and Q_{η} respectively. We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5, P_4 = 13, Q_0 = 5, Q_1 = 2, Q_2 = 6, Q_3 = 17$ and $Q_4 = 46$.

Corollary 5.2. For $\eta \geq 1$, we have the followings:

(a) $\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} P_{-\lambda} = \frac{1}{9}((-1)^{\eta} ((3\eta - 8)P_{-\eta+4} - (9\eta - 21)P_{-\eta+3} + (6\eta - 4)P_{-\eta+2} - (9\eta - 6)P_{-\eta+1} + (6\eta + 11)P_{-\eta}) + 1).$

(b) $\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} Q_{-\lambda} = \frac{1}{9}((-1)^{\eta} ((3\eta - 8)Q_{-\eta+4} - (9\eta - 21)Q_{-\eta+3} + (6\eta - 4)Q_{-\eta+2} - (9\eta - 6)Q_{-\eta+1} + (6\eta + 11)Q_{-\eta}) - 32).$

At this point, recall theorem 4.1 and set $r = s = t = u = 1$ and $v = 2$; this gives the result below.

Proposition 5.3. For $\eta \geq 1$ and setting $r, s, t, u = 1$ and $v = 2$, in theorem 4.1 leads to the next outcomes.

$$\sum_{\lambda=1}^{\eta} \lambda(-1)^{\lambda} A_{-\lambda} = \frac{1}{9}((-1)^{\eta} ((3\eta - 4)A_{-\eta+4} - (6\eta - 5)A_{-\eta+3} + (3\eta + 5)A_{-\eta+2} - (6\eta + 4)A_{-\eta+1} + (3\eta + 14)A_{-\eta}) + 4A_4 - 5A_3 - 5A_2 + 4A_1 - 14A_0).$$

The next corollary follows if we replace A_η in the proposition 5.3 with followings respectively:

- $A_\eta = J_\eta$ with $J_0 = 0, J_1 = 1, J_2 = 1, J_3 = 1, J_4 = 1,$
- $A_\eta = j_\eta$ with $j_0 = 2, j_1 = 1, j_2 = 5, j_3 = 10, j_4 = 20,$
- $A_\eta = K_\eta$ with $K_0 = 3, K_1 = 1, K_2 = 3, K_3 = 10, K_4 = 20,$
- $A_\eta = Q_\eta$ with $Q_0 = 3, Q_1 = 0, Q_2 = 2, Q_3 = 8, Q_4 = 16,$
- $A_\eta = S_\eta$ with $S_0 = 0, S_1 = 1, S_2 = 1, S_3 = 2, S_4 = 4,$
- $A_\eta = R_\eta$ with $R_0 = 5, R_1 = 1, R_2 = 3, R_3 = 7, R_4 = 15.$

Corollary 5.3. For $\eta \geq 1$, we get the followings:

- (a) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda J_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)J_{-\eta+4} - (6\eta - 5)J_{-\eta+3} + (3\eta + 5)J_{-\eta+2} - (6\eta + 4)J_{-\eta+1} + (3\eta + 14)J_{-\eta}) - 2).$
- (b) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda j_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)j_{-\eta+4} - (6\eta - 5)j_{-\eta+3} + (3\eta + 5)j_{-\eta+2} - (6\eta + 4)j_{-\eta+1} + (3\eta + 14)j_{-\eta}) - 19).$
- (c) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda K_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)K_{-\eta+4} - (6\eta - 5)K_{-\eta+3} + (3\eta + 5)K_{-\eta+2} - (6\eta + 4)K_{-\eta+1} + (3\eta + 14)K_{-\eta}) - 23).$
- (d) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda Q_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)Q_{-\eta+4} - (6\eta - 5)Q_{-\eta+3} + (3\eta + 5)Q_{-\eta+2} - (6\eta + 4)Q_{-\eta+1} + (3\eta + 14)Q_{-\eta}) - 28).$
- (e) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda S_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)S_{-\eta+4} - (6\eta - 5)S_{-\eta+3} + (3\eta + 5)S_{-\eta+2} - (6\eta + 4)S_{-\eta+1} + (3\eta + 14)S_{-\eta}) + 5).$
- (f) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda R_{-\lambda} = \frac{1}{9}((-1)^\eta ((3\eta - 4)R_{-\eta+4} - (6\eta - 5)R_{-\eta+3} + (3\eta + 5)R_{-\eta+2} - (6\eta + 4)R_{-\eta+1} + (3\eta + 14)R_{-\eta}) - 56).$

Proposition 5.4. For $\eta \geq 1$ and setting $r = 2, s = 3, t = 5, u = 7, v = 11$, in theorem 4.1 leads to the next outcomes.

$$\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda A_{-\lambda} = \frac{1}{81}((-1)^\eta ((9\eta + 2)A_{-\eta+4} - (27\eta + 15)A_{-\eta+3} + 36A_{-\eta+2} - (45\eta + 46)A_{-\eta+1} - (18\eta - 77)A_{-\eta}) - 2A_4 + 15A_3 - 36A_2 + 46A_1 - 77A_0).$$

The next corollary follows if we replace A_η in the proposition 5.4 with G_η, H_η and E_η respectively. We additionally apply the initial conditions where $G_0 = 0, G_1 = 0, G_2 = 0, G_3 = 1, G_4 = 2, H_0 = 5, H_1 = 2, H_2 = 10, H_3 = 41, H_4 = 150, E_0 = 0, E_1 = 0, E_2 = 0, E_3 = 1$ and $E_4 = 1$.

Corollary 5.4. For $\eta \geq 1$, we get the followings:

- (a) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda G_{-\lambda} = \frac{1}{81}((-1)^\eta ((9\eta + 2)G_{-\eta+4} - (27\eta + 15)G_{-\eta+3} + 36G_{-\eta+2} - (45\eta + 46)G_{-\eta+1} - (18\eta - 77)G_{-\eta}) + 11).$
- (b) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda H_{-\lambda} = \frac{1}{81}((-1)^\eta ((9\eta + 2)H_{-\eta+4} - (27\eta + 15)H_{-\eta+3} + 36H_{-\eta+2} - (45\eta + 46)H_{-\eta+1} - (18\eta - 77)H_{-\eta}) - 338).$
- (c) $\sum_{\lambda=1}^{\eta} \lambda(-1)^\lambda E_{-\lambda} = \frac{1}{81}((-1)^\eta ((9\eta + 2)E_{-\eta+4} - (27\eta + 15)E_{-\eta+3} + 36E_{-\eta+2} - (45\eta + 46)E_{-\eta+1} - (18\eta - 77)E_{-\eta}) + 13).$

5.3 The case $z = i$

In this subsection, we present the closed-form solutions (identities) for sums $\sum_{\lambda=1}^{\eta} \lambda i^\lambda A_{-\lambda}$, specifically considering the sequence $\{A_\eta\}_{\eta \geq 0}$. At this point, recall theorem 4.1 and set $r = s = t = u = v = 1$; this gives the result below.

Proposition 5.5. For $\eta \geq 1$ and setting $r, s, t, u, v = 1$, in theorem 4.1 leads to the next outcomes.

$$\sum_{\lambda=1}^{\eta} \lambda i^{\lambda} A_{-\lambda} = \frac{1}{2i} (i^{\eta} (((1+i)\eta - 6 - i)A_{-\eta+4} + (6 - 2\eta - 4i)A_{-\eta+3} + (10 + 5i - (1 + 3i)\eta)A_{-\eta+2} + ((2 - 2i)\eta + 4 + 10i)A_{-\eta+1} + ((1+i)\eta - 2 + 7i)A_{-\eta}) + (6 + i)A_4 - (6 - 4i)A_3 - (10 + 5i)A_2 - (4 + 10i)A_1 + (2 - 7i)A_0).$$

The next corollary follows if we replace A_{η} in the proposition 5.5 with P_{η} and Q_{η} respectively. We additionally apply the initial conditions where $P_0 = 0, P_1 = 1, P_2 = 2, P_3 = 5, P_4 = 4, Q_0 = 5, Q_1 = 1, Q_2 = 3, Q_3 = 7$ and $Q_4 = 15$.

Corollary 5.5. For $\eta \geq 1$, we obtain the followings.

(a) $\sum_{\lambda=1}^{\eta} \lambda i^{\lambda} P_{-\lambda} = \frac{1}{2i} (i^{\eta} (((1+i)\eta - 6 - i)P_{-\eta+4} + (6 - 2\eta - 4i)P_{-\eta+3} + (10 + 5i - (1 + 3i)\eta)P_{-\eta+2} + ((2 - 2i)\eta + 4 + 10i)P_{-\eta+1} + ((1+i)\eta - 2 + 7i)P_{-\eta}) + (-2 - 3i).$

(b) $\sum_{\lambda=1}^{\eta} \lambda i^{\lambda} Q_{-\lambda} = \frac{1}{2i} (i^{\eta} (((1+i)\eta - 6 - i)Q_{-\eta+4} + (6 - 2\eta - 4i)Q_{-\eta+3} + (10 + 5i - (1 + 3i)\eta)Q_{-\eta+2} + ((2 - 2i)\eta + 4 + 10i)Q_{-\eta+1} + ((1+i)\eta - 2 + 7i)Q_{-\eta}) + (24 - 17i).$

Corresponding summations of the other fifth-order generalized Pentanacci numbers can be calculated in a similar way.

6 Conclusion

In this study, we initially revisited the definition of generalized Pentanacci numbers, laying out the formal expressions and recurrence relations. We also drew from several important publications in the literature to build a foundation for our paper. To enhance comprehension, we present three detailed tables. Table 1 showcases the notations of some special cases of the generalized Pentanacci sequences with the related references. Table 2 gives the notations of some members of generalized Pentanacci sequences with the OEIS numbers of some of them and related papers. Finally, Table 3 presents the some special studies of sum formulas related to sequences and related studies. Moreover, we highlight two crucial theorems conducted by Soykan in [32], which have proven invaluable in this research. These theorems, referenced frequently throughout this paper, provided key theorems that significantly contributed to the development of the results presented here.

In Section 2, we compute the following sums for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with positive subscripts: $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}, \sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$.

In section 3, we provide the closed-form solutions (identities) for sums $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{\lambda}, \sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda}$ and $\sum_{\lambda=0}^{\eta} \lambda z^{\lambda} A_{2\lambda+1}$ for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$. The above sums have been calculated for the sequences listed in Table 2 when $z = -1$ and $z = i$.

In section 4, we compute the following sum for the generalized Pentanacci sequences $\{A_{\eta}\}_{\eta \geq 0}$ with negative subscripts: $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$.

In section 5, we provide the closed-form solutions (identities) for sums $\sum_{\lambda=1}^{\eta} \lambda z^{\lambda} A_{-\lambda}$ for particular cases of $z = -1$ and $z = i$, specifically considering the sequence $\{A_{\eta}\}_{\eta \geq 0}$. The above sums have been computed for the sequences listed in Table 2 when $z = -1$ and $z = i$.

Disclaimer (Artificial Intelligence)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Melham RS. Some Analogs of the Identity $F_{\eta}^2 + F_{\eta+1}^2 = F_{2\eta+1}^2$. *Fibonacci Quarterly*. 1999;37(4):305-311.
- [2] Natividad LR. On solving fibonacci-like sequences of fourth, fifth and sixth order. *International Journal of Mathematics and Computing*. 20133(2), 38-40.
- [3] Rathore GPS, Sikhwal O, Choudhary R.. Formula for finding nth Term of Fibonacci-Like Sequence of Higher Order. *International Journal of Mathematics And its Applications*. 2016;4(2-D):75-80.
- [4] Soykan Y. On generalized pentanacci and gaussian generalized pentanacci numbers. *Asian Research Journal of Mathematics*. 2020;16(9):102-121.
DOI:10.9734/ARJOM/2020/v16i930224
- [5] Soykan Y. Properties of generalized fifth-order pell umbers. *Asian Research Journal of Mathematics*. 2019;15(3):1-18.
- [6] Soykan Y, PolathEE. A note on fifth order jacobsthal numbers. *IOSR Journal of Mathematics (IOSR-JM)*. 2021;17(2):01-23.
DOI: 10.9790/5728-1702010123
- [7] Soykan Y. A study on generalized 5-primes numbers. *Journal of Scientific Perspectives*. 2020;4(3):185-202.
DOI: <https://doi.org/10.26900/jsp.4.017>.
- [8] Sloane NJ. A. The on-line encyclopedia of integer sequences.
Available: <http://oeis.org/>
- [9] Cook CK, Bacon MR. Some identities for Jacobsthal and Jacobsthal-Lucas numbers satisfying higher order recurrence relations. *Annales Mathematicae et Informaticae*. 2013;41:27-39.
- [10] Akbulak M, Öteleş A. On the sum of Pell and Jacobsthal numbers by matrix method. *Bulletin of the Iranian Mathematical Society*. 2014;40(4):1017-1025.
- [11] Gökbaş H, Köse H. Some sum formulas for products of pell and pell-lucas numbers. *Int. J. Adv. Appl. Math. and Mech*. 2017;4(4):1-4.
- [12] Öteleş A, Akbulak M. A note on generalized k-pell numbers and their determinantal representation. *Journal of Analysis and Number Theory*. 2016;4(2):153-158.
- [13] Koshy T. *Fibonacci and Lucas Numbers with Applications*. A Wiley-Interscience Publication, New York; 2001.
- [14] Koshy T. *Pell and pell-lucas numbers with applications*, Springer, New York; 2014
- [15] Hansen RT. General identities for linear fibonacci and lucas summations. *Fibonacci Quarterly*. 1978;16(2):121-28.
- [16] Soykan Y. On Summing Formulas For Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers. *Advances in Research*. 2019;20(2):1-15.

- [17] Soykan Y. Corrigendum: On Summing Formulas for Generalized Fibonacci and Gaussian Generalized Fibonacci Numbers. *Advances in Research*. 2020;21(10):66-82.
DOI: 10.9734/AIR/2020/v21i1030253
- [18] Soykan Y. On Summing Formulas for Horadam Numbers. *Asian Journal of Advanced Research and Reports*. 2020;8(1):45-61.
DOI: 10.9734/AJARR/2020/v8i130192.
- [19] Soykan Y. Generalized fibonacci numbers: Sum Formulas. *Journal of Advances in Mathematics and Computer Science*. 2020;35(1):89-104.
DOI:10.9734/JAMCS/2020/v35i130241.
- [20] Soykan Y. Generalized tribonacci numbers: Summing formulas. *Int. J. Adv. Appl. Math. and Mech*. 2020;7(3):57-76.
- [21] Soykan Y. On Sum Formulas for Generalized Tribonacci Sequence. *Journal of Scientific Research & Reports*. 2020;26(7):27-52.
DOI: 10.9734/JSRR/2020/v26i730283.
- [22] Frontczak R. Sums of Tribonacci and Tribonacci-Lucas Numbers. *International Journal of Mathematical Analysis*. 2018;12(1):19-24.
- [23] Parpar T. k'ncü Mertebeden Rekürans Bağıntısın ın Özellikleri ve Bazı Uygulamaları. Selçuk Üniversitesi, Fen Bilimleri Enstitüsü. Yüksek Lisans Tezi. (In Turkish); 2011.
- [24] Soykan Y. Summing Formulas For Generalized Tribonacci Numbers. *Universal Journal of Mathematics Applications*. 2020;3(1):1-11.
DOI: <https://doi.org/10.32323/ujma.637876>
- [25] Soykan Y. Summation Formulas for Generalized Tetranacci Numbers. *Asian Journal of Advanced Research and Reports*. 2019;7(2):1-12.
DOI:10.9734/ajarr/2019/v7i230170.
- [26] Soykan Y. Matrix sequences of tribonacci and tribonacci-lucas numbers. *Communications in Mathematics and Applications*. 2020;11(2):281-295.
DOI: 10.26713/cma.v11i2.1102
- [27] Waddill ME. The tetranacci sequence and generalizations. *Fibonacci Quarterly*. 1992;30(1):9-20.
- [28] Soykan Y. Sum Formulas For Generalized Fifth-Order Linear Recurrence Sequences. *Journal of Advances in Mathematics and Computer Science*. 2019;34(5):1-14.
Article no. JAMCS.53303, ISSN: 2456-9968.
DOI: 10.9734/JAMCS/2019/v34i530224.
- [29] Soykan Y. Linear summing formulas of generalized pentanacci and gaussian generalized pentanacci numbers. *Journal of Advanced in Mathematics and Computer Science*. 2019;33(3):1-14.
- [30] Soykan Y. On Summing Formulas of Generalized Hexanacci and Gaussian Generalized Hexanacci Numbers. *Asian Research Journal of Mathematics*. 2019;14(4):1-14. Article no.ARJOM.50727.
- [31] Soykan Y. A study on sum formulas of generalized sixth-order linear recurrence sequences. *Asian Journal of Advanced Research and Reports*. 2020;14(2):36-48.
DOI:10.9734/AJARR/2020/v14i230329

- [32] Soykan Y. A study on sum formulas of generalized pentanacci sequence: Closed forms of the sum formulas $\sum_{k=0}^{\eta} x^k W_k$ and $\sum_{k=1}^{\eta} x^k W_{-k}$. Journal of Progressive Research in Mathematics. 2021;18(2):20-38.
- [33] Soykan Y. Sum formulas of generalized pentanacci numbers: Closed forms of the sum formulas $\sum_{k=0}^{\eta} k W_k$ and $\sum_{k=1}^{\eta} k W_{-k}$, Int. J. Adv. Appl. Math. and Mech. 2021;8(4):1-14.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/124484>