



Gastrointestinal Helminths Parasites of Cattle Slaughtered in Gwagwalada Abattoirs

Balarabe-Musa Binta ^a, Onyeagba Kosisochukwu Dematus ^a and Airuoyuwa Peace ^{a*}

^a Department of Biological Sciences, Faculty of Science, University of Abuja, P.M.B 117, Federal Capital Territory, Abuja, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Somdet Srichairatanakool, Chiang Mai University, Thailand.

Reviewers:

(1) Maria Priscila Saracino, Universidad de Buenos Aires, Argentina.

(2) Attahiru Adamu, Gombe State College of Nursing and Midwifery, Nigeria.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/78531>

Original Research Article

Received 11 October 2021

Accepted 22 December 2021

Published 29 December 2021

ABSTRACT

A survey was conducted to determine the prevalence of gastrointestinal helminth parasites in cattle slaughtered in Gwagwalada abattoirs, and a total of 165 gastrointestinal helminth parasites belonging to four genera were reported. *Ascaris lumbricoides*, *Fasciola gigantica*, *Taenia saginata*, and *Trichuris trichiura* are the four genera and four species of helminth parasites found. The abattoir at Angwandodo recorded twenty-seven (27) *Ascaris lumbricoides* intestinal helminth parasites, six (6) *Fasciola gigantica* parasites, and three (3) *Taenia saginata* parasites. Similarly, (21) *Taenia saginata* was identified, followed by (12) *Ascaris lumbricoides*, and (9) *Trichuris trichiura* and *Fasciola gigantica* were reported at the Gwako abattoir. Furthermore, the abattoir in Giri was home to (18) *Ascaris lumbricoides*, (15) *Taenia saginata*, twelve (12) *Trichuris trichiura*, and three (3) *Fasciola gigantica*. *Taenia saginata* and *Ascaris lumbricoides* each recorded nine (9) samples, while *Trichuris trichiura* and *Fasciola gigantica* each recorded six (6) samples from an abattoir in Kutunku. From the total of one hundred and sixty-five (165) gastrointestinal parasites observed in cattle slaughtered in abattoirs, sixty-six (66) were *Ascaris lumbricoides*, fifty-one (51) were *Taenia saginata*, twenty-seven (27) were *Trichuris trichiura* (16.36 percent), and twenty-one (21) were *Fasciola gigantica*, the least parasites observed with a 12.73 percent prevalence. There was a significant difference in the prevalence of gastrointestinal helminths in cattle slaughtered in

*Corresponding author: E-mail: onyeagbakosi001@gmail.com;

Gwagwalada abattoirs ($p < 0.05$). The livestock industry plays a vital role in the economy of Nigeria. It serves as a major source of income and livelihood for majority of Nigerians who are rural settlers and contributes about 5.2% to the National Gross Domestic Product (GDP). Helminths however, cause economic losses due to reductions in milk production, weight gain, fertility and carcass quality. Zoonotic helminths of livestock origin cause health problems in humans. Hence, there is a need to monitor intestinal parasites of cattle to promote animal production and public health in Nigeria.

Keywords: Gastrointestinal helminths; Ascaris lumbricoides; prevalence; parasites.

1. INTRODUCTION

Cattle, Nigeria's most common domesticated livestock, are an asset in both traditional and modern agriculture. Furthermore, they provide meat, milk, skin, and draught power for farming [1]. In Nigeria, the livestock sector accounts for 5.2 percent of the GDP, while cattle production accounts for 50 percent of total meat production [2]. Helminths are well-known to be a major impediment to ruminant well-being and productivity [3,4]. Gastrointestinal helminths are common parasitic agents of livestock, particularly ruminants, and are known to limit cattle production in a variety of areas and countries [5].

At times, the mortality of animals due to parasitic diseases may not be alarming, but their indirect effects on livestock productivity and the way they are passed from animals to humans are considerably greater [6]. Indirect losses associated with helminth infections include the reduction in productive potential, such as decreased growth rate, weight loss, diarrhea, anorexia, and sometimes anemia [7,8] discovered that grazing habits, climate, nutritional deficiency, pasture management, immunological status, vector, presence of intermediate host, and the number of infective larvae and eggs in the environment are the most important predisposing factors for helminth infections. The effect of helminth infections is determined by a combination of factors, of which the varying susceptibility of the host species, the pathogenicity of the parasite species, the host and parasite interaction, and the infective dose are the most important [9]. However, helminth zoonotic diseases do not only involve animals but are also considered a major health problem in many countries. There is a need to carry out this study on the intestinal helminths parasites of cattle slaughtered in Gwagwalada to know the prevalence of the helminthic parasites responsible for cattle diseases in Gwagwalada.

2. MATERIALS AND METHODS

2.1 Study Region

This study was carried out at the Biology laboratory of the Department of Biological Sciences at the University of Abuja, Gwagwalada, FCT-Abuja, Nigeria. Gwagwalada is one of the five municipal councils of the Federal Capital Territory of Nigeria, together with Abaji, Kuje, Bwari, and Kwali. The FCT also includes the City of Abuja. Gwagwalada is also the name of the main town in the Local Government Area, which has an area of 1,043 km² and a population of 157,770 at the 2016 census and is located between latitudes 8° 55' and 8° 60' North and 7° 05' and 7° 11' East [10]. The Gwagwalada abattoirs are located in different areas, which include Agwandodo, Gwako, Giri, and the New Kutunku ward of the town, beside one of the tributary streams of the river Usuma, which drains through the town. It has a recorded annual temperature that ranges from 30 degrees Celsius to 37 degrees Celsius and a total annual rainfall of about 1650 mm. Gwagwalada abattoir is located in a high-density residential area and consists of three sections; the slaughtering sections, processing sections, and dumping sections.

2.2 Study Population

At Gwagwalada, FCT Abuja abattoirs, the butchers only slaughter adult male cattle; therefore, the research was based only on male cattle.

2.3 Sample Collections

A total of 160 fecal samples of freshly slaughtered cattle were randomly collected from the rectum, with 40 samples each from four (4) different abattoirs located in Gwagwalada; FCT Abuja. Fecal samples were collected from the abattoirs at Kutunku, Agwandodo, Gwako, and Giri with five (5) samples per week for eight (8)

weeks. The samples were collected in sterile plastic containers and taken to the Biological Sciences laboratory at the University of Abuja, Gwagwalada FCT-Abuja- Nigeria, for the study of helminthic parasites in the intestinal tract of the cattle. The samples were analyzed on the day of collection.

2.4 Evaluation of Helminthic Parasites

The fecal sample was spread on a dissecting board and visible worms to the naked eye were picked up using thumb forceps and then stored in 10% formalin before identification. About one gram (1 g) of the fecal sample was dissolved in a sterile test tube containing 10 mL of sterile distilled water. The mixtures were agitated for 3 min by shaking and then centrifuged at 4000 rpm for 10 min. The supernatant was decanted and the sediment was examined for helminthic parasites [11]. Microscopic examination of the helminthic parasites was done at X40 magnification. The parasites were identified based on their shape, size, and color.

2.5 Statistical Analysis

The prevalence of intestinal helminths parasites was statistically analyzed using one-way ANOVA and the F-test statistic from Microsoft Excel Statistics at a P = 0.05 level of significance.

3. RESULTS

3.1 Prevalence of Gastrointestinal Helminth Parasites in Slaughtered Cattle

Table 1 shows the prevalence of gastrointestinal helminth parasites in cattle slaughtered in abattoirs in the Gwagwalada Area Council. Four genera and four species of helminth parasites were observed and included *Ascaris lumbricoides*, *Fasciola gigantica*, *Taenia saginata*, and *Trichuris trichiura*, respectively, as seen in Table 1. Twenty-seven (27) of the intestinal helminth parasites were *Ascaris lumbricoides* and *Fasciola gigantica*, six (6), while three (3) *Taenia saginata* were recorded for the abattoir in Agwandodo. Similarly, twenty-one (21) *Taenia saginata* was recorded, followed by twelve (12) *Ascaris lumbricoides*, which were also recorded, while *Trichuris trichiura* and *Fasciola gigantica* were nine (9) each, for the abattoir in Gwako. Furthermore, eighteen (18) *Ascaris lumbricoides* were recorded, followed by fifteen (15) *Taenia saginata* and *Trichuris*

trichiura, and only three (3) *Fasciola gigantica* were recorded for the abattoir in Giri. Also, *Taenia saginata* and *Ascaris lumbricoides* recorded nine (9) each, while *Trichuris trichiura* and *Fasciola gigantica* recorded six (6) each, for samples collected from an abattoir in Kutunku, as seen in Table 1.

3.1.1 Frequency and percentage of gastrointestinal helminth parasites in cattle slaughtered in abattoirs

From overall total of one hundred and sixty-five (165) gastrointestinal helminth parasite observed in cattle slaughtered in the abattoirs, sixty-six (66), equivalent to 40 % of the total number of parasites observed which was the highest number, were *Ascaris lumbricoides* and fifty-one (51) of them which is equivalent to 30.91 % were *Taenia saginata* while twenty-seven (27) were *Trichuris trichiura* (16.36 %) and only twenty-one were *Fasciola gigantica* being the least parasite observed with 12.73 % as seen in Table 2.

From the above Table, the test statistic is the F value of 5.46 and at 0.05 level of significant, the critical $F_{0.05; 4,15} = 3.59$. Since the test statistic is higher than the critical value, the null hypothesis (H_0) is rejected and the alternative hypothesis is accepted meaning that there is significant difference between the prevalence of intestinal helminth parasites in cattle slaughtered in abattoirs in Gwagwalada.

4. DISCUSSION

The findings of this study show that all the cattle screened had helminth infection, thus providing valuable information on the burden of helminths among cattle in Nigeria, since animals slaughtered in this abattoir are representative of cattle in the country. The overall prevalence of helminth infection obtained in this study is similar to that of Edosomwan and Shoyemi (2012), who reported a prevalence of 167.1% in the south-south region of Nigeria but higher than the 50.8% and 62.1% earlier reported by [11] in south-eastern and south-southern Nigeria, respectively. The differences observed could be due to the periods or seasons in which the studies were conducted as well as the sources of cattle sampled in the various regions. The most prevalent helminth parasite observed in cattle slaughtered was *Ascaris lumbricoides* with sixty-six (66), equivalent to 40% of the total number of parasites observed, and fifty-one (51) of them, which is equivalent to 30.91% were *Taenia saginata*, while twenty-seven (27) were *Trichuris*

trichiura (16.36%), and only twenty-one were *Fasciola gigantica*, the least parasite observed with 12.73% as seen in Table 2. [12] also reported the prevalence of *Taenia* sp, *Ascaris* sp; and *Trichuris trichiura* in livestock slaughtered in Dei-Dei Abattoir, F.C.T Abuja. Furthermore, this finding revealed that nematodes and cestodes were the most prevalent among the helminths. However, this is at variance with previous reports by [13,14,15] reported nematodes as the most prevalent helminths in studies carried out in India, Ethiopia, and eastern Nigeria, respectively. This difference could, however, be associated with differences in geographical and/or climatic conditions and ecology. Again, the helminths identified in this study were similar to those identified by [16,15] in earlier studies carried out in Benin and Port Harcourt, south-south Nigeria. It can therefore be suggested that the similarity in the helminth profile indicates exposure of these animals to common conditions (e.g., ecology, pasture, and humidity) which are prevalent in northern Nigeria, where most of these animals

are sourced from before being transported to different abattoirs in Nigeria. One major factor that would have accounted for this is the fact that cattle in the local setting in Nigeria are exposed to poor feeding and veterinary care, factors accountable for equal susceptibility to helminth infections. The difference in the prevalence obtained could be attributed to the existence of favorable environmental factors necessary for the prolonged survival and development of the infective larval stage of most helminths. Furthermore, the management system of animals could also be accountable for the difference in prevalence. Cattle with a moderate body condition score had a higher prevalence of intestinal helminths when compared to those that were emaciated. Possible explanations for this include the fact that those with moderate body condition tolerated helminth infections better for a variety of reasons, including good nutrition, or that both host and parasite had reached a state of equilibrium and were asymptomatic at the time of sample collection.

Table 1. Frequency of gastrointestinal helminth parasites in cattle slaughtered in Gwagwalada abattoirs

Abattoirs locations	Number of Sample	ParasitesSpecies	Frequencies of Occurrence
Agwandodo	40	<i>Fasciola gigantica</i>	3
		<i>Taenia saginata</i>	6
		<i>Ascaris lumbricoides</i>	27
Gwako	40	<i>Taenia saginata</i>	21
		<i>Trichuris trichiura</i>	9
		<i>Fasciola gigantica</i>	9
		<i>Ascaris lumbricoides</i>	12
Giri	40	<i>Taenia saginata</i>	15
		<i>Trichuris trichiura</i>	12
		<i>Fasciola gigantica</i>	3
		<i>Ascaris lumbricoides</i>	18
Kutunku	40	<i>Taenia saginata</i>	9
		<i>Trichuris trichiura</i>	6
		<i>Fasciola gigantica</i>	6
		<i>Ascaris lumbricoides</i>	9
Total	160		165

Table 2. Percentage of occurrence of intestinal helminth parasites in cattle slaughtered in abattoirs in Gwagwalada Area Council

Parasite species	Frequencies	Percentages (%)
<i>Ascaris lumbricoides</i>	66	40
<i>Taenia saginata</i>	51	30.91
<i>Trichuris trichiura</i>	27	16.36
<i>Fasciola gigantica</i>	21	12.73
Total	165	100

Table 3. Summary

Groups	Count	Sum	Average	Variance
<i>Trichuris</i> sp	3	9	3	1
<i>Taenia</i> sp	4	17	4.25	4.92
<i>Toxocara</i> sp	4	7	1.75	0.92
<i>Ascaris</i> sp	4	26	6.5	4.33

Table 4. Anova value

Source of Variation	SS	Df	MS	F	P-value	F critical
Between Groups	48.43	3	16.14	5.46	0.02	3.59




Fig. 1. Total intestinal helminth parasites in cattle slaughtered different abattoirs in Gwagwalada




Fig. 2. Overall Prevalence of intestinal helminth parasite in cattle slaughtered different abattoirs in Gwagwalada

5. CONCLUSION

The result of this study showed a moderately high prevalence of intestinal helminth infection of both economic and zoonotic importance among trade cattle slaughtered in abattoirs in Gwagwalada, Nigeria. This has a negative impact on both animal production and public health. Therefore, to mitigate these problems, appropriate anthelmintic regimens and control measures (i.e., comprehensive parasite control programs, pasture management, and environmental sanitation) in cattle and public health awareness should be encouraged. Finally, there is a need to monitor intestinal parasites in cattle to promote animal production and public health in Nigeria.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

ACKNOWLEDGEMENTS

The authors thank all those who have been associated with this project, especially the staff of the department of Biological Sciences at the University of Abuja and the personnel at the abattoirs, for their cooperation throughout the period of this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Adedipe NO, Bakshi JS, Odegbaro OA, Aliyu A. Study on gastrointestinal helminth infections in small-scale dairy cattle farms of Jimma town Ethiopia. *Journal of Applied Sciences and Technology*. 2016; 34: 45-65.
2. Awowole B, Francis. This is a waste. *Daily sun*; The sun publishing limited, Lagos. *New Journal*. 2017;10-23.
3. Dantanko H, Idris HS. Helminthosis in Livestock Slaughtered in Dei-Dei Abattoir. *F.C.T Abuja. Global Advanced Research Journal of Agricultural Science*. 2014;3(9):304-309.
4. Edosomwan EU, Shoyemi OO. Prevalence of gastrointestinal helminth parasites of cattle and goats slaughtered at abattoirs in Benin City, Nigeria. *African Scientist*. 2012;13(2):109-114.
5. Ekong PS, Juryit R, Dika NM, Nguku P, Musenero M. Prevalence and risk factors for zoonotic helminth infection among humans and animals, Jos, Nigeria. *The Pan African Medical Journal*. 2012; 11: 22-6.
6. Elele K, Owhoeli O, Gboeloh LB. Prevalence of species of helminths parasites in cattle slaughtered in selected abattoirs in Port Harcourt, south-south, Nigeria. *International Research on Medical Sciences*. 2013;1(2):10-17.
7. Food and Agricultural Organisation. Distribution and impact of helminth diseases of livestock in developing countries, in *FAO Corporate Document. Repository Agriculture and Consumer Protection*. 2011; 22: 45-66.
8. Haliu T, Cherenet A, Moti Y, Tadele T. Study on gastrointestinal helminth infections in small-scale dairy cattle farms of Jimma town, Ethiopia. *Ethiopian Journal of Applied Sciences and Technology*. 2011;2(1):31-37.
9. Hesterberg UW, Bagnall R, Perrett K, Horner R, Gummow B. A questionnaire survey of perceptions and preventive measures related to animal health amongst cattle owners of rural communities in KwaZulu-Natal, South Africa. *Journal of the South African Veterinary Association*. 2017;78(4):205-208.
10. Keyyu JD, Monrad J, Kyvsgaard NC, Kassuku AA. Epidemiology of *Fasciola gigantica* and amphistomes in cattle on traditional, small-scale dairy and large-scale dairy farms in the southern highlands of Tanzania. *Tropical Animal Health and Production*. 2015;37(4):303-314.
11. Mir MR, Chishti MZ, Rashid Z. Point prevalence of gastrointestinal helminthiasis in large ruminants of Jammu, India. *International Journal of Scientific and Research Publications*. 2013;3(8):2013.

12. Radostits OM, Blood DC, Gay CC. "Diseases caused by helminth parasites, in Veterinary Medicine: A Textbook of Diseases of Cattle, Sheep, Pigs, Goats and Horses. BailliereTindall, London, UK, Journal 8th edition. 2014;1223–1230.
13. Rajput ZI, Hu SW, Chen J, Arijo AG. Xiao CW. Importance of ticks and their chemical and immunological control in livestock. Science Journal of Zhejiang University. 2016;7(11):912–921.
14. Swai ES, Mtui PF, Mbise AN, Kaaya E, Sanka P, Loomu PM. Prevalence of gastro intestinal parasite infections in Maasai cattle in Ngorongoro District, Tanzania. Livestock Research for Rural Development. 2016;18: 8.
15. Tewe OO. Sustainability and Development": Paradigms from Nigeria's Livestock Industry, Inaugural Lecture series. University of Ibadan Press, Ibadan, Nigeria. 2017;97.
16. Nwigwe J, Njoku OO. Odikamnor, O.O and Uhuo, C.A. Comparative study of intestinal helminths and protozoa of cattle and goats in Abakaliki metropolis of Ebonyi State, Nigeria. Advances in Applied Science Research. 2013;4(2):223–227.

© 2021 Binta et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78531>