



Hematemia Revealing Cytomegalovirus Esophagitis in an Immunocompetent Patient: A Case Report

Mouna Tamzaourte^{1*}, Ahlam Benhamdane¹, Sanae Berrag¹, Tarik Adioui¹ and Aziz Aourarh¹

¹*Gastroenterology I unit, Mohamed V Military Teaching Hospital, Mohamed V-Souissi University, Rabat, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. P. Veera Muthumari, V.V.Vanniaperumal College for Women, India.

Reviewers:

(1) Maysa de Oliveira Barbosa, Cariri Regional University, Brazil.

(2) Mohammad Fadhil Ibraheem, University of Baghdad, Iraq.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/78357>

Received 01 October 2021

Accepted 04 December 2021

Published 09 December 2021

Case Study

ABSTRACT

Cytomegalovirus (CMV) esophagitis is well-documented in immunocompromised patients. A few studies have described CMV infection in immunocompetent patients diagnosed with a critical illness. However, CMV esophagitis has rarely been documented in immunocompetent hosts. We report a case of CMV esophagitis in an immunocompetent patient who presented with gastrointestinal bleeding that was successfully resolved with valganciclovir treatment. Our case supports observations that CMV esophagitis can cause significant morbidity, regardless of immune system status.

Keywords: Cytomegalovirus; esophagitis; immunocompetent.

1. INTRODUCTION

Cytomegalovirus (CMV) is a member of the Herpesviridae family. Although immunocompetent individuals can be infected,

the serious disease typically only manifests in those with immunocompromised states. CMV can affect many organs, including the eyes, stomach, colon, but the focus of this discussion is on the esophagus. We reported a case of

*Corresponding author: Email: mouna.tamzaourte@gmail.com;

hematemia revealing cytomegalovirus esophagitis in an immunocompetent patient and to highlight the diagnostic and therapeutic aspects of this rare entity.

2. PRESENTATION OF CASE

A 60 years old man with a history of diabetes complicated with retinopathy; high blood pressure and former smoker. He had been taking insulin therapy and amlodipin. He underwent eso-gastroduodenal fibroscopy for melena in 2019, which revealed an ulcer located in the anterior surface of the bulbe treated with proton pump inhibitor. He also underwent right colectomy for colon cancer in January 2020.

The patient was admitted to the emergency department with hematemesis and melena. He also complained of palpitations and dizziness. The patient's vital signs were stable with blood pressure: 130/80 mmhg; heart rate: 82 beats/min; respiration rate: 16/min; and body temperature 37°C. The physical examination revealed a chronically ill-looking man with median laparotomy scar. Laboratory analysis showed normochromic normocytic anemia (Hb=5,6 g/dl), and leukocytosis (10400 cells/mL) with no other abnormalities. He was transfused with 3 red blood cells with control hemoglobin 9,2g/dl. Eso-gastroduodenal fibroscopy revealed severe ulcerative esophagitis and ulcerated bulbo-duodenitis with recent bleeding marks. Biopsy specimens from the esophageal ulcer showed an ulcerated mucosa with an intense polymorphic inflammatory infiltrate. Immunohistochemical staining using the anti-CMV monoclonal antibody was positive, while the patient showed negative results for CMV antigenemia. The CMV Polymerase Chain Reaction (PCR) on esophageal biopsy was also positive. The serology test was negative for IgM anti CMV with a titer of 0,3 U/ml but positive for IgG anti CMV with a titer of 250U/ml. Complementary study with syphilis screening, serology for human immunodeficiency virus 1 et 2, hepatitis B and C viruses were negative. Serology for herpes simplex virus 1 revealed previous infection (IgM negative and IgG positive) and Epstein Barr virus serology was negative. The diagnostic of CMV ulcerative esophagitis was made. The patient was treated with oral valganciclovir 450 mg 2 times a day for 15 days as well as treatment with proton pump inhibitor type omeprazole at a dose of 8 mg per hour by electric syringe pump in the acute phase for 48 hours then 80 mg per day intravenously

then orally for 8 weeks. The outcome was favorable with cessation of bleeding and clinical improvement. Endoscopic control carried out 2 months later was normal with total disappearance of the esophageal lesions.

3. DISCUSSION

Cytomegalovirus is a common infectious pathogen and the majority of cases are mild symptomatic or asymptomatic [1]. It is a member of the Herpesviridae family of viruses which has the largest genome of any herpes viruses [2].

CMV is often inactive and may exist in a latent stage before becoming reactivated. It may be acquired during the perinatal period or in adulthood through sexual contact [3].

CMV infection is generally infrequent in immunocompetent patients. Although those can be infected, severe disease typically only manifests in immunocompromised status such as AIDS patients, organ transplant recipients and patients on chemotherapy [4]. CMV can affect almost any area of the gastrointestinal tract. CMV esophagitis is the second most common gastrointestinal tract manifestation of the disease, the first being colitis [5]. The most common symptoms of CMV esophagitis are dysphagia, odynophagia and epigastric pain [6]. Other non specific symptoms include nausea, vomiting, abdominal pain, weight loss and diarrhea [5]. Hematemesis is a rare manifestation of CMV esophagitis compared with CMV gastritis or duodenitis [7, 8]. The endoscopic appearance of CMV esophagitis consisted of multiple well demarcated, vertical or horizontal, superficial linear ulcers at the median to the distal esophagus [9]. Ulcerations can be single or multiple, can be deep resembling cavitation, and have associated features such as mucosal edema and nodularity [10].

There is no pathognomonic endoscopic feature for CMV esophagitis, and definitive diagnosis depends heavily on biopsy findings. Histopathologic diagnosis can be made by identifying CMV inclusion bodies with standard H and E staining, called a cytomegalic cell [3, 11]. The advent of immunohistochemical staining via anti CMV monoclonal antibodies has provided more modalities to help aid in the diagnosis, reaching a sensitivity of 78% to 93% [12].

To date, the mainstay of CMV esophagitis treatment is intravenous Ganciclovir or oral

Valganciclovir [13]. In case of resistance to the first line treatment, Foscarnet is an alternative to Ganciclovir. Failure of monotherapy requires an attempt at combined therapy with IV Ganciclovir and IV Foscarnet [3].

4. CONCLUSION

Although hematemesis and esophageal ulcers are rare, they should be considered as potential complications accompanying CMV infection in immunocompetent individuals. The administration of symptomatic and antiviral therapy should be considered even when the patient is immunocompetent.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bate SL, Dollard SC, Cannon MJ. Cytomegalovirus seroprevalence in the United States: the national health and nutrition examination surveys, 1988-2004. *Clin Infect Dis*. 2010 Jun 1;50(11):1439-47. DOI: 10.1086/652438.
2. Isaacson MK, Compton T. Human Cytomegalovirus glycoprotein B is required for virus entry and cell-to-cell spread but not for virion attachment, assembly, or egress. *J Virol* 2009 Apr;83(8):3891-903. DOI: 10.1128/JVI.01251-08. Epub 2009 Feb 4.
3. Goodgame RW. Gastrointestinal cytomegalovirus disease. *Ann Intern Med*. 1993 Nov 1;119(9):924-35. DOI:10.7326/0003-4819-119-9-199311010-00010.
4. Li L, Chakinala RC. Cytomegalovirus Esophagitis. [Updated 2021 May 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-

Available: <https://www.ncbi.nlm.nih.gov/books/NBK556110/>

5. Wang HW, Kuo CJ, Lin WR, HSU CM, HO YP, Lin CJ et al. The clinical characteristics and manifestations of cytomegalovirus esophagitis. *Dis Esophagus*. 2016 May;29(4):392-9. DOI: 10.1111/dote.12340. Epub 2015 Feb 26
6. Shumpei Y, Masaya I, Muneaki M, Naoyuki N, Motowo M, Hiroyuki O. Severe Bleeding due to Cytomegalovirus Esophagitis in a Patient with Diabetes after Interbody Fusion Surgery. *Intern Med*. 2009 Oct 15;58(20):2949-2955. DOI:10.2169/internalmedicine.2908-19
7. Peter A, Telkes G, Varga M, Sarvary E, Kovalszky I. Endoscopic diagnosis of cytomegalovirus infection of upper gastrointestinal tract in solid organ transplant recipients: Hungarian single center experience. *Clin Transplant*. 2004 Oct;18(5):580-4.
8. Featherstone RJ, Camero LG, Khatib R, Snower D, Mungara P. Massive esophageal bleeding in achalasia complicated by cytomegalovirus esophagitis. *Ann Thorac Surg* 1995 Apr;59(4):1021-2. DOI: 10.1016/0003-4975(94)00744-r. PMID: 7695381.
9. Khetpal R, Arnaoutakis K, Dholoria B. CMV esophagitis: sequelae of radiation therapy. *Esophagus* 2013;10:244-246
10. Marques S, Carmo J, Pinto D, Bispo M, Ramos S, Chagas C. Cytomegalovirus Disease of the upper Gastrointestinal tract: A 10 year retrospective study. *GE Port J Gastroenterol*. 2017 Nov;24(6):262-268. DOI: 10.1159/000479232. Epub 2017 Sep 2. PMID: 29255766; PMCID: PMC5731150.
11. Juric-Sekhar G, Upton MP, Swanson PE, Westerhoff M. Cytomegalovirus (CMV) in gastrointestinal mucosal biopsies: should a pathologist perform CMV immunohistochemistry if the clinician requests it? *Hum Pathol*. 2017 Feb;60:11-15. DOI: 10.1016/j.humpath.2016.09.009. Epub 2016 Sep 22. PMID: 27666768.
12. Gravito-Soares E, Gravito-Soares M, Camacho E, Tome L. Cytomegalovirus ulcerative esophagitis in a young healthy immunocompetent patient. *BMJ Case Rep*. 2018 Mar 5;2018:bcr2017223297. DOI: 10.1136/bcr-2017-223297. PMID: 29507025; PMCID: PMC5847831.

13. Emery VC. Cytomegalovirus: recent progress in understanding pathogenesis and control. QJM. 2012 May;105 (5): 401-405. DOI: 10.1093/qjmed/hcr262. Epub 2011 Dec 22. PMID: 22198913; PMCID: PMC3328974.

© 2021 Tamzaourte et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78357>*