



Edaphic Factors and Flooding Periodicity Determining Forest Types in a Topographic Gradient in the Northern Brazilian Amazonia

**L. A. Beltrán Alemán^{1*}, R. I. Barbosa², I. Montero Fernández³,
L. C. S. Carvalho^{1,4}, P. E. Barni⁵, R. L. C. Oliveira⁶, J. F. Vale Jr.⁷,
S. A. Saravia Maldonado⁸ and N. E. Arriaga Pérez⁸**

¹Post-Graduate Program in Natural Resources, PRONAT, Federal University of Roraima, Campus Paricarana, Boa Vista, RR, Brazil.

²Department of Environmental Dynamics, National Institute for Research in Amazonia (INPA), Rua Coronel Pinto 315, 69301-150 Boa Vista, Roraima, Brazil.

³Postgraduate Program in Biodiversity and Biotechnology - Roraima, Federal University of Roraima, Campus Paricarana, Boa Vista, RR, Brazil.

⁴University of Exeter, Prince of Wales Road, Exeter, Devon, EX4 4SB, UK.

⁵Universidade Estadual de Roraima (UERR), Campus Rorainópolis, Av. Senador Helio Campos, s/n, 69375-000 Rorainópolis, Roraima, Brazil.

⁶Universidade Estadual de Roraima, Coordenação do Curso de Ciências Biológicas, Rua Sete de Setembro, 231, Canarinho, Boa Vista, 69306-530, Roraima, Brazil.

⁷Departamento de Solos e Engenharia Agrícola, Universidade Federal de Roraima, Campus Cauamé, BR 174, Km 12, Bairro Monte Cristo, CEP 69300-000, Boa Vista, Roraima, Brazil.

⁸National University of Agriculture, Highway to Dulce Nombre de Culmí, Neighborhood El Espino Catacamas-Olancho, Honduras.

Authors' contributions

This work was carried out in collaboration among all the authors. The authors LABA, RIB, IMF, LCSC, PEB, RLCO and JFVJ determined the area of study and the methodology to be used in the field and in the drafting of the protocol and the first draft of the manuscript. The authors LABA and RIB managed the study analyzes. The authors LABA, RIB, IMF, LCSC, SASM and NEAP assisted in the literature search and review of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2019/v28i630127

Editor(s):

(1) Dr. Surendra Singh Bargali, Professor, Department of Botany, DSB Campus, Kumaun University, Nainital, Uttarakhand, India.

Reviewers:

(1) Donald DeAngelis, United States.

(2) Sergio Esteban Lozano Baez, University of São Paulo, Brazil.

(3) Junir Antonio Lutinski, Unochapecó, Brazil.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50026>

Original Research Article

**Received 02 May 2019
Accepted 10 July 2019
Published 18 July 2019**

ABSTRACT

The Brazilian Amazonia is a region covered by an extensive mosaic of tropical forests conditioned by different topographical and hydro-edaphic features. Although studies relating environmental determinants of structure and floristic composition are systematically evolving in the region, there is no doubt that there are still information gaps due to the lack of research in peripheral areas of the Amazonia. The seasonally flooded areas of the state of Roraima situated on rio Branco-rio Negro basin, northern Brazilian Amazonia, still are deprived of such information. In this way, this work had as objective determine the physical and soil chemical attributes, and the flooding periodicity that characterize different forest types dispersed in a topographic gradient located in an area on the north of rio Branco-rio Negro basin. Soil samples (0-60 cm) were collected along a 2.7 km transect (31.1-64.8 m a.s.l.) crossing three different forest types: (i) mosaic between treed and forested shade-loving (La+Ld), (ii) area of ecological tension between forested shade-loving and open ombrophilous forest (LO) and (iii) open ombrophilous forest (Ab+As). The results indicated different soil classes and flooding periodicity for each forest type observed: Entisols Fluvents (La+Ld, 3-4 months flooded), Entisols Quartzipsamments (LO, 1-2 months) and Yellow Ultisols (Ab+As, no flooding). All analyzed soils were defined as nutrient-poor areas, especially those located on low altitude, characterized for higher hydrological restrictions (seasonal flooding) aggregating forest types of lower structural patterns (e.g. La+Ld). Soils on low altitude were also characterized as those with the highest percentage of fine sand and silt, while soil free of seasonal flooding (Yellow Ultisols) presented the highest levels of clay and coarse sand, always associated with the ombrophilous forests (higher structural patterns). These results improve our understanding of the environmental factors conditioning different forest types in this peripheral region of Amazonia, suggesting that ecosystems with higher hydro-edaphic restrictions are a strong indicator of forest types with lower structural patterns.

Keywords: Oligotrophic ecosystems; water table; ecotone; phytophysiology.

1. INTRODUCTION

The Amazon basin occupies ~40% of the surface of South America and about 60% is inserted within the Brazilian territory [1]. Its predominant covering is defined as dense and open ombrophilous forests [2-3], but throughout the Amazonian biome there are many different forest types that may be distinguished by their floristic composition and structure due to the large environmental heterogeneity [4-5]. In general, the factors modeling the different forest types are attributed mainly to the climatic variations, hydro-edaphic conditions, topography, and anthropogenic interferences, all interacting and acting at different spatial scales [6-9]. This congregation of factors generates different structural and floristic shades with different ecosystem values, but generally information on the weight of each one, in the local and regional context, is little known due to the gigantism of the Amazonia, which makes sampling in peripheral areas a difficult process [3,10]. However, this kind of information is the basis for improving our knowledge on specialization of the various Amazonian phytophysionomies, with different structural patterns and species diversity, directly

influencing the estimates of biomass/carbon stocks and fluxes in this region considered to be the largest and most important "natural environment" mitigating the harmful effects of global climate change [11-12].

Although studies involving edaphic and hydrological factors in association with topographic gradients related to the dynamics of ecosystems have evolved rapidly in the Amazon [14-16], there is still a great lack of information due to huge regional gaps with rare scientific investigations. In this context, ecotone forests (transition areas, contact zones or areas of ecological tension) are ecoregions representing about 15% of the biome [17] but still have a lack of knowledge about the processes of their formation and maintenance [18]. These ecoregions are characterized by mosaics of different forest types condensed into distinct spatial scales that hamper their floristic and structural characterization and, above all, biomass/carbon estimates [19,20]. This scarcity is mainly detected in the northern of the Brazilian Amazonia, especially in the seasonally flooded areas of the rio Branco-rio Negro basin located in the state of Roraima [16,21-22].

Recent works have demonstrated that the ecosystems which form the mosaic of landscapes in this region present a direct integration between the plant cover and the physical, chemical and biological attributes of the soil [23] due to essential processes related to the biogeochemical cycles, water Table outcrops, accumulation and decomposition of organic matter [16,24]. This is a strong indication that environmental conditions associated with temporal flooding, sediment drag, and nutrient leaching may have high importance in the formatting of different forest types in this peripheral region of the Amazonia [25-27]. In the same context, the physical and chemical attributes of the soil, altitude, flooding periodicity, drainage and microclimate also considered as important determinants in the formation of different natural environments with specific structural patterns [28-29].

The objective of this study was to determine the flooding periodicity and the physical and soil chemical attributes that characterize a topographic gradient established between the Água Boa do Univini River and the Cumaru Mountain, a peripheral area of the Rio Branco-Rio Negro basin. This region belongs to the Serra da Mocidade National Park, a federal protected area located in the state of Roraima, northern Brazilian Amazonia. This region is formed by a large mosaic of forest and non-forest ecosystems without rare scientific investigations about the role of hydro-edaphic factors as determinants of different phytophysiognomies formations. Our results aim to improve the understanding of the environmental factors that determine different forest types in this region of Amazonia, indicating the association between environments with higher/lower hydro-edaphic restrictions and their respective forest structural pattern taking into account horizontal (stem diameter) and vertical (total height) parameters.

2. MATERIALS AND METHODS

2.1 Study Area

This study was carried out in the Serra da Mocidade National Park (1.405°N - 61.648°W and 1.382°N - 61.673°W), a federal protected area managed by ICMBio (Chico Mendes Institute for Biodiversity and Conservation), located in the municipality of Caracará, ~ 290 km south of Boa Vista, capital of Roraima. The sampling area is characterized by ecotone zones (area of ecological tension or transition area or

contact zone) between seasonally flooded forests under strong influence of the Água Nova do Univini river (black water) in association with open ombrophilous forests that reach the first steps of elevation of the Cumaru Mountain (Fig. 1). The phytophysiological characterization, edaphic and flood periodicity were performed in a topographic gradient (31.1-64.8 m a.s.l.) inserted in an irregular transect (2.7 km long) known as "Cat's Trail", which begins on the right bank of the Água Boa do Univini river to the first steps of the northeast sector of the Cumaru Mountain, located at the northeastern end of the National Park (Fig. 1).

2.1.1 Characterization of the study area

The National Park is totally inserted in the rainy tropical monsoon type (Am) following Köppen classification, with annual rainfall of 1700-2000 mm and May-August representing the rainiest period (~40% of annual rainfall [30-31]. This region is marked by a chain of mountains that lends its name (Serra da Mocidade), resulting from the erosion of the Guyanese Craton, a large continental block formed by magmatic and metamorphic rocks dated between 1.8-2.5 billion years, in the Lower Pre-Cambrian period [32]. The characteristics of the main soil types in the National Park region are defined from lithological residues of the same geological constitution of the rock formation complex of the Serra da Mocidade, being a large residual mass, with an altitude reaching ~1.800 m, characterized by sharp crests and ravine slopes, covered by high altitude forests that lose this characteristic when reaching the zones of flooded forests of low altitude and smaller biometric structural patterns. Along this rocky complex, eight different soil classes can be found: Neossolo Flúvico (Entisols Fluvents), Neossolo Quartzarênico (Entisols Quartzipsamments), Espodossolo Humilúvico (Spodosols), Latossolo Amarelo (Yellow Oxisols), Gleissolo Háplico (Entsols), Latossolo Vermelho (Red Oxisols), Neossolo Litólico (Entsols Lithic) and Argissolo Amarelo (Yellow Ultisols) [32-33].

2.2 Phyto-characterization and Periodicity of Flooding

In order to carry out the phytophysionomic classification of the forest types in the sample area, we adopted the criteria proposed by the Brazilian Vegetation Classification System [34], based on a forest inventory carried out along the entire transect at Cat's Trail (R. I. Barbosa,




Fig. 1. Study area indicating the geographical location of the Serra da Mocidade National Park and the soil profiles sampled along the Cat's Trail

personal communication). In this survey, the structure (horizontal and vertical) and arboreal groups (trees and palms) with stem diameter ≥ 10 cm were defined in each forest type arranged along the topographic gradient (31,1-64, 8 m a.s.l.). The flood periodicity data were obtained from observations performed in two consecutive rainy periods (2016 and 2017), where the sampling transect was coursed from start to finish in both periods, estimating a mean time interval (months) of flooding for each of them. All information was aggregated into an ecosystem conceptual model that faithfully followed the observed topographic gradient. Later, this model was adopted as an associative basis of the chemical and soil physical characteristics under each defined forest type.

2.3 Soil Sampling and Physical/Chemical Analyzes

In order to analyze and describe the physical and chemical attributes of the soil, three profiles (1 m wide, 1 m long, 80 cm deep) were opened for each forest type considered. Soil samples were collected at 0-20 cm, 20-40 cm and 40-60 cm depths. After that, the samples were deposited in plastic bags and identified by forest type and depth. All samples were air dried (TFSA), sieved (2 mm) and sent to the Soil Laboratory for physical (% sand, % silt and % clay) and chemical (pH, organic matter, exchangeable acidity, potential acidity, Ca, Mg, K, P, Cu, Zn, Fe, Mn and B) analysis following the methodology specified [35]. The descriptive classification of

the soils sampled was performed by the Brazilian Soil Classification System [36] up to the third categorical level and correlated to the Soil Taxonomy [33]. The authorization number issued by the Chico Mendes Institute for the Conservation of Biodiversity for the conduct of this study was: (Authorization SISBEn, 54221-1).

3. RESULTS AND DISCUSSION

The three profiles open along the sampled transect of the Serra da Mocidade National Park delimited three different types of soils: (i) Entisols Fluvents, (ii) Entisols Quartzipsamments and (iii) Yellow Ultisols (Fig. 2). All of them were characterized as chemically poor (Table 1) and with high sand contents (Table 2). The results found in the forest type classified as Treed and Forested shade-loving (La + Ld), 3-4 month flooded in the year, are associated with the Entisols Fluvents, while Entisols Quartzipsamments and Yellow Ultisols are associated, respectively, to the Area of Ecological Tension (LO) and Open Ombrophylous Forest (Ab+As) (Table 3). These hydro-edaphic characteristics are similar to those reported in the Viruá National Park [21-22], a region with similar ecological characteristics to the Serra da Mocidade, strongly indicating that soils with higher levels of sand, low nutrient contents and higher flooding periodicity can define oligotrophic forest types with lower structural patterns (vertical and horizontal), in counterpoint to areas free of flooding processes, such as ombrophilous forests (Fig. 3).

Table 1. Chemical attributes determined in three profiles along a topographic gradient located in the Serra da Mocidade National Park, in the north of the Brazilian Amazonia in 2018

Depth (cm)	pH H ₂ O	OM g/kg	P	Cu	Fe	Zn	Mn	B	K	Ca	Mg	H+Al	Al	SB	CEC	Sat.	Sat.
																Bases V%	Al m%
Entisols Fluvents– Treed and Forested shade-loving (La+Ld)																	
0-20	5.1	14	3	6.80	92.00	8.20	3.00	0.43	0.10	0.20	0.10	3.40	0.20	0.40	3.80	11	33
20-40	5.2	7	2	6.70	37.20	8.05	1.40	0.25	0.06	0.10	0.10	2.80	0.20	0.26	3.06	9	43
40-60	4.8	5	2	4.60	14.20	5.80	1.10	0.24	0.06	0.10	0.10	2.50	0.20	0.26	2.76	9	43
Entisols Quartzipsamments – Area of Ecological Tension (LO)																	
0-20	4.6	23	3	1.60	9.80	5.65	15.40	0.26	0.20	0.30	0.10	6.40	0.40	0.60	7.00	9	40
20-40	4.9	9	3	2.10	18.60	3.80	2.40	0.38	0.15	0.30	0.10	4.20	0.30	0.55	4.75	12	35
40-60	4.9	5	2	2.00	19.00	3.60	1.20	0.35	0.13	0.30	0.10	3.80	0.20	0.53	4.33	12	27
Yellow Ultisols – Open Ombrophylous Forest (Ab+As)																	
0-20	4.6	11	3	2.90	106.00	5.95	6.50	0.34	0.15	0.60	0.20	4.70	0.20	0.95	5.65	17	17
20-40	4.7	5	2	2.10	102.00	4.30	1.90	0.32	0.12	0.20	0.10	3.40	0.30	0.42	3.82	11	42
40-60	4.6	5	1	0.70	80.00	3.45	2.00	0.27	0.06	0.10	0.10	3.40	0.20	0.26	3.66	7	43

Where: OM = organic matter, P = Phosphorus, Cu = Copper, Fe = Iron, Zn = Zinc, Mn = Manganese, B = Boron, K = Potassium, Ca = Calcium, Mg = Magnesium, H + Al = Acidable exchangeable, Al = Aluminium exchangeable, SB = Sum of bases, CEC =Cationic Exchange Capacity, V = Saturation by Bases, m =Saturation by Aluminium



Fig. 2. Soil profiles opened in the Serra da Mocidade National Park
 Where the profiles area: A= Entisols Fluvents (Treed and Forested shade-loving La+Ld) B= Entisols Quartzipsamments (Area of Ecological Tension LO) and C= Yellow Ultisols (Open Ombrophyllous Forest Ab+As)




Fig. 3. Conceptual model of the three forest types characterized by the soil classes and flooding periodicity in a transect in the Serra da Mocidade National Park, northern Brazilian Amazonia

Entisols Fluvents can be considered as a low developed soil formed by quaternary sediment deposits, where drainage varies from moderate to imperfect, and it is very influenced by the outflow of the water table [37]. In the study area, this soil class was characterized by the predominance of high levels of fine sand followed by the silt (Table 2), associated with mosaics of treed and forested shade-loving, which are oligotrophic forest environments with low structural expression that naturally dominate part of the rio Branco-rio Negro basin [38-40]. The pH in this environment showed a tendency to reduce as a function of soil depth, due to the high levels of fine sand, which facilitates the processes of water percolation and nutrient leaching [41]. The available phosphorus also presented a decrease with depth, being this a process associated to alluvial sediment entrainment condition of the high parts of the topographic gradient, considering that the parts of lower altitude are characterized by more restrictive soils (3-4 months flooded) that may eventually accumulate

higher levels of fertility in the early layers in contrast with deeper layers [26].

Our sampling in the profile of this soil class indicated that the organic matter had the highest values in the first sampling layer (0-20 cm). This result is in agreement with that presented [25-42], where the authors suggest that soils with higher elevations tend to have higher clay contents compared to lowland soils, but the organic matter contents are higher in the first layers of lowland soils due to the strong drag provided by the topography of terrain. Concerning to the physical properties of this soil class situated in the La+Ld forest type, it was verified higher concentrations of silt and fine sand, which are textural particles fully compatible with natural lowland environment [43].

Entisols Quartzipsamments is characterized as a soil under Area of Ecological Tension (LO), 1-2 months flooded in the year, and it can be described as a soil class extremely weathered, much quartzous, where almost all the clay is

Table 2. Physical attributes determined in three profiles along a topographic gradient located in the Serra da Mocidade National Park, northern Brazilian Amazonia in 2018

Depth (cm)	Arg	Sil	ArT	ArG	ArF
%				
Entisols Fluvents – Treed and Forested shade-loving (La+Ld)					
0-20	13.9	19.1	67.0	7.0	60.0
20-40	14.2	16.8	69.0	4.0	65.0
40-60	14.2	16.8	69.0	4.0	65.0
Entisols Quartzipsamments – Area of Ecological Tension (LO)					
0-20	15.7	11.3	73.0	20.0	53.0
20-40	17.4	9.6	73.0	21.0	52.0
40-60	16.9	8.1	75.0	23.0	52.0
Yellow Ultisols – Open Ombrophylous Forest (Ab+As)					
0-20	15.4	3.6	81.0	51.0	30.0
20-40	20.3	6.7	73.0	42.0	31.0
40-60	25.0	6.0	69.0	37.0	32.0

Where: Arg Clay (<0.002mm), Sil Silt (0.053-0.002mm), Total ArT_Area, ArG_ Thick Sand (2.00-0.210mm), ArF_ Fine Sand (0.210-0.053mm)

Table 3. Soil classes, flood periodicity and structural parameters (DBH and total height) of the forest types observed in the sampling transect formed by the Cat’s Trail, Serra da Mocidade National Park. Where: La + Ld = Mosaic of treed and forested shade-loving; LO = Area of ecological tension between forested shade-loving and ombrophylous forest; Ab + As = open ombrophylous forest associated with the first steps of the Cumarú Mountain. Different uppercase (Trees, ANOVA followed by Tukey test) and lowercase (Palms; Test t) letters in the columns indicate discrepancies ($\alpha = 0.05$) between the values of the taxonomic groups

Forest Type	Density (ind ha ⁻¹)		DBH (cm)		Ht (m)	
	Trees	Palms	Trees	Palms	Trees	Palms
La+Ld	940	0	13.7± 3.3 A	-	12.2±2.3 A	-
LO	710	45	17.3± 7.4 B	14.5±3.5 a	17.7±2.7 B	17.5±1.9 a
Ab+As	423	85	20.9±12.9 C	18.5±6.5 b	18.3±4.0 B	16.6±3.2 a

(*) DBH = diameter at breast height (cm) and Ht = total height (m)

destroyed by acidolysis, or sandy deposits formed by wind phenomena, occurring in flat reliefs or basin reliefs or even in soft undulating reliefs [36-37]. This class of soil presented high levels of fine and coarse sand (Table 2), where these physical characteristics are associated to the formation processes of this environment, especially by the presence of small streams that precede the first undulating steps of the Cumarú Mountain, with undulations which aid the accumulation of alluvial material derived from the highest part. Thus, this soil type is related to the hydrological processes of sediment trapping, as also observed in the region of the National Park of Viruá (Roraima), in similar environments to those found in Serra da Mocidade [21].

The pH values determined for this soil class in the study area are in line with the standards [35]. An analysis along the soil profile in this forest type allowed to understand that there is a slight increase in the pH values from the most

superficial layer (0-20 cm) to the deepest ones (20-40 cm, 40-60 cm) and, consequently, a reduction in exchangeable and potential acidity (Table 1). This is a process fed by the infiltration of exchangeable bases or increase of organic matter, as previously established [44]. Likewise, the values of available phosphorus presented a decrease between the second (20-40 cm) and the third layer (40-60 cm), being a strong indicative of the reduction of this element along the vertical soil profile, as observed [45] and it can act as a limiting element in the larger / smaller vertical and horizontal structuring of the forest.

Organic matter also declines from the superficial to the deeper layers of this soil class in the study area. This same observation was reported in the Viruá National Park, with the authors suggesting that the topography of the terrain, especially those with soft ripples, may retain organic matter in the superficial layers of the soil due to the

sediment trapping or the temporal outcropping of the water table [21] which are the same environmental characteristics observed in the ecotone (LO) of the Cat's Trail. In the same sense of the organic matter, the CEC also has a reduction pattern from the most superficial to the deepest layers, presenting a CEC saturation problem with the exchangeable Al, being common in this soil type, which can be aggravated by depth due to the decrease of organic matter in the soil and exchangeable bases such as Ca^{2+} , Mg^{2+} and K^+ , which would limit the development of the roots of the plants and affect the structure of the forest [46].

The Yellow Ultisols is characterized as a soil type flooding free, with strong presence of Open Ombrophylous Forest (Ab+As), with high contents of coarse sand and clay (Table 2) the formation factors of this soil class are similar to that of the Oxisols, with the same geomorphological characteristics and natural vegetation, but with a textural gradient [37]. The pH values found in this environment are slightly acidic and there is not much difference between the layers (Table 1). This process may be happening due to the absence of both water table outcrops and temporal flooding in this area [41-47]. The content of organic matter found in this soil type was high in the first layer, decreasing drastically towards the deeper layers. This result indicates a direct relationship with the CEC values found in this soil type, within the relational congruence [48] where organic matter and clays are the edaphic parameters with the greatest contribution to the formation of CEC values.

In this same analytical path, the concentrations of the micronutrients observed in the profile of this soil highlight the higher concentration of Fe in the first two layers (0-20 cm and 20-40 cm), being this element the main constituent of the structure of the clays [49]. This chemical characteristic was also observed in the ombrophilous forests of the Viruá National Park [23] suggesting that higher Fe contents in the ombrophilous forests of regions under the influence of treed and forested shade-loving mosaics may be due to the high presence of mineral particles (oxides of iron) derived from the organic matter deposited on the soil in litter form. This indicates that both Fe and the other micronutrients (Zn, Mn, B, Cu) have an important role in the nutrient cycling of this forest environment, but without a clearly defined role as a characterizer of forest types. This evidence the

large range of uncertainties that still persist in the evaluations on the relationships between hydro-edaphic conditions and their role in the construction of Amazonian ecosystems. In this way, it is inferred that a better spatial distribution of the pedo-phytosociological studies, addressing peripheral regional gaps, can help us to generate environmental standards (topographic gradients, periodicity of flooding, soil classes) that more accurately configure the particular structural and floristic characteristics of different forest types, allowing better accuracy in biomass/carbon estimates, and giving the real importance of the Amazonia in the context of the global warming mitigation.

4. CONCLUSION

We conclude that each soil class determined in this study has a strong association with the topographic gradient sampled in the study area situated on Serra da Mocidade National Park, where lower altitude environments with larger flooding periods are related to forest types of lower structural pattern (e.g. treed and forested shade-loving) preferentially on oligotrophic soils (poor and sandy). These characteristics indicate the formation of environments influenced by continuous hydro-edaphic and geological processes, where seasonal flooding and sediment trawling are part of the process of formation of the main forest types in the study area. Therefore, edaphic factors and flooding periodicity are environmental characteristics that act as environmental filters which are important in the formation of the landscape in this region of rio Branco-rio Negro basin. These results improve our understanding of the environmental factors that determine different forest types in this region of the Amazonia, where environments with higher hydro-edaphic restrictions are a strong indicator of forest types with lower vertical and horizontal structure.

ACKNOWLEDGMENT

This study was supported for two institutional projects: (i) Ecologia e Manejo dos Recursos Naturais de Savanas e Florestas de Roraima (PPI-INPA 015/122) and (ii) Instituto Nacional de Ciência e Tecnologia dos Serviços Ambientais da Amazônia (INCT ServAmb-Processo FAPEAM/CNPq). Luis Antonio Beltrán Alemán was supported by postgraduate fellowships from the Organization of American States (OAS), R. I. Barbosa by a research grant from Conselho Nacional de Pesquisas (CNPq Proc.

304204/2015-3) and L.C.S. Carvalho by a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES/PNPD). The ICMBio team, Érika Tieko Fujisaki, Sylvio Romério Briglia-Ferreira and Inara Rocha, provided both support in all trips to the Serra da Mocidade National Park and authorization for the study (Authorization SISBIO n. 54221-1). The topographer Neptali Portillo Villanueva contributed formatting all figures of the study area.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Carvalho ST, Domingues PE. Economic and deforestation scenario for the Brazilian Amazon between 2006 and 2030, *Nova Economia*. 2016;1(26):585-621.
2. Veloso HP, Rangel Filho ALR, Lima JCA. Classificação da vegetação brasileira adaptada a um sistema universal. Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro. 1991;124.
3. Fearnside PM. Quantificação do serviço ambiental do carbono nas florestas amazônicas brasileiras. *Oecologia Brasiliensis*. 2008;12(4):743-756.
4. Fearnside PM, Ferraz J. A conservation gap analysis of Brazil's Amazonian vegetation. *Conservation Biology*. 1995; 9(5):1134-1147.
5. Nogueira EM, Yanai AM, Fonseca FO, Fearnside PM. Carbon stock loss from deforestation through 2013 in Brazilian Amazonia. *Global Change Biology*. 2015; 21:1271–1292.
6. Castilho CV, Magnusson WE, Araújo RNO, Luizão RCC, Luizão FJ, Lima AP, Higuchi N. Variation in aboveground tree live biomass in a central Amazonian Forest: Effects of soil and topography. *Forest Ecology and Management*. 2006;234(1-3): 85-96.
7. Phillips OP, Vargas O, Monteagudo AL, Cruz AP, Zans M, Sánchez W, Yli-halla M, Rose, R. Habitat association among Amazonian tree species: A landscape-scale approach. *Journal of Ecology*. 2003; 91(1):757-775.
8. Laurance WF, Fearnside PM, Laurance, SG, Delamonica P, Lovejoy TE, Rankin-de-Merona JM, Chambers JQ, Gascon C. Relationship between soils and Amazon forest biomass: A landscape-scale study. *Forest Ecology and Management*. 1999; 18:127-138.
9. Franco-Moraes J, Juliano AFB, Baniwa Flávia RC, Costa Helena P, Lima Charles R, Clement Glenn H, Shepard Jr. Historical landscape domestication in ancestral forests with nutrient-poor soils in north-western Amazonia. *FORECO*; 2019.
10. Phillips OL. Changes in the carbon balance of tropical forest: Evidence from long-term plots. *Science*. 1998;282(5388): 439-442.
11. Lewis SL, Phillips OL, Baker TR, Lloyd J, Malhi Y, Almeida S. Concerted changes in tropical forest structure and dynamics: Evidence from 50 South American long-term plots. *Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences*. 2004;359(1443):421-436.
12. IPCC (Intergovernmental Panel on Climate Change). Guidelines for national greenhouse gas inventories. National greenhouse gas inventories programme. Eggleston HS, L. Miwa T, Ngara and K Tanabe (eds). Intergovernmental Panel on Climate Change (IPCC), Institute for Global Environmental Strategies (IGES), JAPAN; 2006.
13. ter Steege H, Vaessen RW, Cárdenas-López D, Sabatier D, Antonelli A, Oliveira SM, Pitman NCA, Jørgensen PM, Salomão RP. The discovery of the Amazonian tree flora with an updated checklist of all known tree taxa. *Scientific reports*. 2016;6:295-349.
14. Tuomisto H, Ruokolainen K, Aguilar M, Sarmiento A. Floristic patterns along a 43-km long transect in an Amazonian rain forest. *Journal of Ecology*. 2003;91:743–756.
15. Tuomisto H, Zuquim G, Cardenas G. Species richness and diversity along edaphic and climatic gradients in Amazonia, *Ecography*. 2014;37(11):1034-1046.
16. Silva LFSG, Castilho CV, Cavalcante CO, Pimentel TP, Fearnside PM, Barbosa RI. Production and stock of coarse woody debris across a hydro-edaphic gradient of oligotrophic forests in the northern Brazilian Amazon. *Forest Ecology and Management*. 2016;364:1-9.
17. Santos CPF, Valle G, Sestini F, Hoffman F, Dousseau L, Homem de Mello J.

- Mapeamento dos remanescentes e ocupação antrópica no Bioma Amazônia; 2007.
Available:<http://mar.te.dpi.inpe.br/rep/dpi.inpe.br/sbsr@80/2006/11.18.01.25?mirror=dpi.inpe.br>
[Acesso em 02 dez. 2017]
18. Santos NMC, Vale Júnior JF, Barbosa RI. Florística e estrutura arbórea de ilhas de mata em áreas de savana do norte da Amazônia brasileira. *Boletim do Museu Paraense Emílio Goeldi (Ciências Naturais)*. 2013;8(2):205-221.
 19. Nascimento M, Carvalho L, Barbosa R, Villeda D. Variation in floristic composition, demography and above-ground biomass over a 20 year period in Amazonian Monodominant forest. *Plant Ecology & Biodiversity*. 2014;7:293-303.
 20. Barni PE, Manzi AO, Condé TM, Barbosa RI, Fearnside PM. Spatial distribution of forest biomass in Brazil's state of Roraima, northern Amazonia. *Forest Ecology and Management*. 2016;377:170-181.
 21. Mendonça BA, Fernandes Filho EI, Schaefer CEGR, Simas FNB, do Vale Junior JF, Lisboa B de AR, de Mendonça JGF. Solos e Geoambientes do Parque Nacional do Viruá e entorno, Roraima: visão integrada da paisagem e serviço ambiental. *Ciência Florestal*. 2013;23:427-442.
 22. Damasco G, Vicentini A, Castilho CV, Pimentel TP, Nascimento HEM. Disentangling the role of edaphic variability, flooding regime and topography of Amazonian white-sand vegetation. *Journal of Vegetation Science*. 2013;24(2): 384–394.
 23. Mendonça BAF, Fernandes EIF, Schaefer C, Mendonça JGF, Vasconcelos BNF. Soil-vegetation relationships and community structure in a "terra-firme"-white-sand vegetation gradient in Viruá National Park, northern Amazon, Brazil. *Anais da Academia Brasileira de Ciências*. 2017;89(2):1269-1293.
 24. Cordeiro C, Rossetti DF, Gribel R, Tuomisto H. Impact of sedimentary processes on white-sand vegetation in an Amazonian megafan. *Journal of Tropical Ecology*. 2016;32(6):498-509.
 25. Luizão FJ, Luizão RCC, Proctor J. Soil acidity and nutrient deficiency in central Amazonian heath forest soils. *Plant Ecol*. 2007;192:209-224.
 26. Suwa R, Sakai T, Santos J, Silva P, Kajimoto T, Ishizuka M, Higuchi N. Meaning of the topographic gradient in stem diameter-Height allometry for precise biomass Estimation of a tropical humid forest in the central Amazon. *JARQ*. 2013;47:109-114.
 27. Junk JW, Wittmann F, Schongart J, Piedade, MA. Classification of the major habitats of Amazonian black-water river floodplains and a comparison with their white-water counterparts, *Wetlands Ecol Manage*. 2015;23:677-693.
 28. Khorramdel S, Koocheki A, Nassiri M, RezaKhorasani M, Evaluation of carbon sequestration potential in corn fields with different management systems. *Soil & Tillage Research*. 2013;133:25-31.
 29. Scaranello, M.A da S, Alves LF, Vieira SA, de Camargo PB, Joly CA, Martinelli LA. Height-diameter relationships of tropical Atlantic moist forest trees in southeastern Brazil. *Scientia Agricola*. 2012;69:26-37.
 30. Barbosa RI. Distribuição das chuvas em Roraima. In Barbosa RI, Ferreira EJJ, Castellón EG (Eds.). *Homem, Ambiente e Ecologia no Estado de Roraima*. INPA-AM, Manaus (AM). 1997;325-335.
 31. Alvares CA, Stape JL, Sentelhas PS, Gonçalves PL, Sparovek G. Koppen's climate classification map for Brazil, *Meteorologische Zeitschrift*. 2014;22:711-728.
 32. Brazil-MME. Projeto RADAMBRASIL - Levantamento de Recursos Naturais. Ministério das Minas e Energia, Departamento Nacional de Produção Mineral, Rio de Janeiro. 1975;475. Available:<http://biblioteca.ibge.gov.br/visualizacao/livros/liv24025.pdf>
 33. USDA. *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys*; 1999.
 34. Brazil-MME. Projeto RADAMBRASIL - Levantamento de Recursos Naturais. Ministério das Minas e Energia, Departamento Nacional de Produção Mineral, Rio de Janeiro. 1975;475. Available:<http://biblioteca.ibge.gov.br/visualizacao/livros/liv24025.pdf>
 35. EMBRAPA. *Manual de Análises Químicas de Solos, Plantas e Fertilizantes*. 2009; 624.
 36. Santos HG. *Sistema Brasileiro de Classificação de Solos*. Embrapa Solos. Brazil. 2018;590.

37. Vale Jr JF, Schaefer CEGR. Solos sob Savanas de Roraima. Gênese, classificação e relações ambientes. Gráfica Ioris: Boa Vista-Roraima. Brazil. 2010;219.
38. Prance GT, Schubart HOR. Notes on the vegetation of Amazonia I. A preliminary note on the origin of the open white sand Campinas of the Lower Rio Negro. Brittonia. 1977;30(1):60-63.
39. Barbosa RI, Ferreira CAC. Biomassa acima do solo de um ecossistema de "campina" em Roraima, norte da Amazônia Brasileira. Acta Amazonica. 2004;34(4): 577-586.
40. Mendonça BAF, Fernandes Filho EI, Schaefer CEGR, Carvalho AF, Vale Jr JF, Corrêa GR. Use of geophysical methods for the study of sandy soils under Campinarana at the National Park of Viruá, Roraima state, Brazilian Amazonia. Journal of Soils and Sediments. 2014; 14(3):525-537.
41. Quesada CA, Lloyd J, Anderson LO, Fyllas NM, Schwarz M, Czimexik CI. Soils of Amazonia with particular reference to the Rain for sites. Biogeosciences Discussion. 2009;6:3851-3921.
42. Ferraz J, (Colocar todos os autores do trabalho). Distribuição dos solos ao longo de dois transectos em floresta primária ao Norte de Manaus (AM). In: Higuchi N, et al. Pesquisas florestais para a conservação da floresta e reabilitação de áreas degradadas da Amazônia. Instituto Nacional de Pesquisas da Amazônia. 1998;111-143.
43. Magalhães RC, Gomes RCM. Mineralogy and chemistry of the lowland soil and its sensibilities in the process of lands falls in community Divino Espírito Santo (Amazonas, Brazil). Soc. & Nat. 2013;25: 609-621.
44. Santos HG, Carvalho Junior W, Dart RO, Aglio MLD, Sousa JS, Pares JG, Fontana A, Martins AL, Oliveira Ap. O novo mapa de solos do Brasil: legenda atualizada. Rio de Janeiro: Embrapa Solos. 2011;67.
45. Duivenvoorden JF. Patterns of tree species richness in rain forests of the middle Caqueta area, Colombia, NW Amazonia. Biotropica. 1996;28:142-158.
46. Sacramento P, Rodrigues LRA, da Cruz MCP, Lugão SMB, de Campos FP, Centurion JF, Ferreira ME. Atributos químicos e físicos de um argissolo cultivado com Panicum maximum Jacq. Cv. IPR-86 Milênio, sob lotação rotacionada e adubado com nitrogênio. R. Bras. Ci. Solo. 2008;32(1):183-193.
47. Scopel I, Peixinho DM, Souza MS, Mariano ZF, da Assunção HF. Formação de areais e perspectivas de uso e manejo de Neossolos Quartzarênicos em Serranópolis (GO). Boletim Goiano de Geografia. 2005;25:11-27.
48. Ostertag R. effects of nitrogen and phosphorus availability on fine root dynamics in Hawaiian montane forests. Ecology. 2001;82:485-499.
49. Tsui CC, Chen ZS, Hsieh CF. Relationships between soil properties and slope position in a lowland rain forest of Southern Taiwan. Geoderma. 2004;123: 131-142.

© 2019 Alemán et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50026>