



Acute Seizure Following Isotretinoin Administration: A Detailed Case Report

May Alsane^{a++*}, Adnan Hajjiah^{a#} and Alasmar El-Shably^{a+}

^a Department of Clinical Pharmacy, Ahmadi Hospital, Kuwait.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2023/v25i2601

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96641>

Case Study

Received: 19/12/2022
Accepted: 21/02/2023
Published: 07/04/2023

ABSTRACT

Isotretinoin is considered a cutting edge treatment for Acne vulgaris. It is licensed for treating severe (cystic) acne vulgaris. Our case report represents a 28 year old female who experienced a rare neurological side-effect after ingesting Isotretinoin.

Keywords: Isotretinoin; retinoids; neurological side-effect; acute seizure.

1. INTRODUCTION

Isotretinoin is derived from a group of drugs called Retinoids, it is primarily used to treat acne. The magnitude of the acne crisis extends psychologically to reflect on self-esteem and confidence. Although Isotretinoin offered a solution to a worldwide problem, it has also

caused other issues that could have otherwise been avoided. This case report sheds a light on one of the adverse effects of Isotretinoin.

2. CASE PRESENTATION

A 28 year old female with no previous medical history of note came to the Emergency Room

⁺⁺ B.Pharm MSc;

[#] M.Pharm MSc;

^{*} M.Sc MD MRCP FRCP;

*Corresponding author: E-mail: msane@kockw.com;

(ER) in Ahmadi Hospital following an episode of generalized tonic clonic seizure. The patient described having blurred vision for a few seconds while taking a walk with her mother at the mall. According to the patient's mother the episode continued for about 10-15 minutes and resolved spontaneously. The first aid service in the mall checked both blood sugar and blood pressure which were normal. The patient did not recall the incident postictal, she continued to have episodes of blurred vision the following week before she attended the ER.

The patient denied having any fever, incontinence, or tongue biting. She also denied prolonged screen time on electronic gadgets. When examined she did not complain of cough, shortness of breath or palpitation. The patient was fully conscious, alert and oriented through the examination process and lying comfortably in bed. The patient was vitally stable, afebrile, and maintaining saturation on room air. Neurologically, power and sensations were maintained, her Glasgow Coma Scale (GCS) was 15/15, gait was normal, and there were no cerebral signs.

Computed Tomography was ordered and revealed no evidence of intra / extra-axial hemorrhage, acute infarction or midline shift. Brain parenchyma showed normal grey white matter differentiation.

MRI Cervical spine was performed using standard protocol on 3 T scanner without IV contrast. No previous studies were available for comparison. It was an unremarkable study. MRI brain and Venogram were also requested and were unremarkable and there were no previous results available for comparison. Patient refused Lumbar Puncture testing (LP) and Electroencephalogram (EEG) were planned in outpatient setting. Results showed no evidence of Epilepsy.

All necessary investigations were performed and found no reasonable explanation as to the patient's chief complaint. The patient admitted ingesting Isotretinoin (30mg once daily) for a duration of eight months. The Naranjo scale was used to assess the casual relationship between the medication and the adverse event, which gave a score of 8.

3. DISCUSSION

The mechanism in which Isotretinoin can cause seizures is not fully understood, however

manufacturers link Isotretinoin with dizziness, drowsiness, and blurred vision as part of the rare side-effects as well as mood changes and aggressive behavior. Some studies suggested that isotretinoin can cause swelling of the brain which in turn will lead to seizure, coma, and death. Patients reported headaches which may be linked to pseudo tumor cerebri; an idiopathic increase in the intracranial pressure leading to swelling and seizure [1,2]. Another explanation might be Biotinidase deficiency; an enzyme responsible for the availability of biotin, which in turn is linked with anti-epileptic medication [1]. Isotretinoin can lead to Biotinidase deficiency the effects of which can lead to seizures as well as ataxia, encephalopathy, alopecia, skin rash, conjunctivitis, IQ and developmental delay [1]. Serious repercussion can lead to irreversible neurological damage and death.

Recent case reports provide evidence of a link between Isotretinoin and seizure onset. A patient experienced a generalized seizure episode four days after receiving 80 mg/day dose of isotretinoin [3]. The episode resolved spontaneously within an hour and was followed by another seizure that involved prolonged loss of consciousness [3]. After regaining consciousness the patient continued to have impaired vision and headache [3]. All investigations were normal and symptoms resolved following abrupt discontinuation of Isotretinoin [3]. Muscles stiffness and superimposed painful spasms have been experienced by another adolescent upon receiving isotretinoin for ten days [4]. The symptoms resolved after administration of diazepam and discontinuation of isotretinoin [4]. Chronic vitamin A intoxication has been described in adults who receive 15 mg/daily for prolonged periods. Such intoxication is linked to several side-effects including headaches and diplopia due to increased intracranial pressure [5]. Isotretinoin-induced psychotic episode was experienced by a teenager after ingesting 30mg/day isotretinoin for three months. Recovery involved administration of Olanzapine and discontinuation of isotretinoin [6]. A cohort study revealed that patients exposed isotretinoin were significantly more at risk of developing a neurological disorder than those on other antibiotics, topical acne medications, or are on non-prescription anti-acne medication [7].

4. CONCLUSION

Although Isotretinoin is not void of adverse reactions. Caution and proper counseling are

advised when prescribing Isotretinoin. Further studies are required to assess the neurological side-effects and provide a definite explanation.

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bremner JD. Isotretinoin and neuropsychiatric side effects: Continued vigilance is needed. *Journal of Affective Disorders Reports*. 2021;6. DOI:<https://doi.org/10.1016/j.jadr.2021.100230>
2. Fahad S, Siddiqui DB. Potential neurologic side effects of oral isotretinoin: A case of peripheral neuropathy. *SKIN The Journal of Cutaneous Medicine*. 2022;6(4). DOI: 10.25251/skin.6.4.13
3. Marroni MBG. Isotretinoin: Possible cause of acute seizure and confusion. *The Annals of Pharmacotherapy*. 1993;27(6): 793-794. DOI: 10.1177/106002809302700626
4. Chroni E, Monastirli A, Tsambaos D. Neuromuscular adverse effects associated with systemic retinoid dermatotherapy. *Drug-Safety*. 2010;33:25–34. Available:<https://doi.org/10.2165/11319020-000000000-00000>
5. Kontaxakis VP, Skourides D, Ferentinos P, et al. Isotretinoin and psychopathology: A review. *Ann Gen Psychiatry*. 2009;8(2). Available:<https://doi.org/10.1186/1744-859X-8-2>
6. Isotretinoin. *Reactions Weekly*. 2020; 1814(196). Available:<https://doi.org/10.1007/s40278-020-81183-8>
7. Paljarvi T, McPherson T, Luciano S, Herttua K, Fazel S. Isotretinoin and adverse neuropsychiatric outcomes: Retrospective cohort study using routine data. *Br J Dermatol*. 2022;187(1):64-72. DOI: 10.1111/bjd.21049. Epub 2022 Apr 12. PMID: 35174880; PMCID: PMC9543533.

© 2023 Alsane et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/96641>*